ELSEVIER
SAUNDERS

Prim Care Clin Office Pract
32 (2005) 1131–1172

PRIMARY CARE:
CLINICS IN
OFFICE PRACTICE

Cumulative Index 2005

Note: Page numbers of article titles are in **boldface** type.

A

Abciximab, for coronary artery disease, 1068

Abdominal wall, posterior, groin pain in, 178–180

Abductor pollicis longus tendon, De Quervain's tenosynovitis of, 37–38, 95

Absorbent products, for urinary incontinence, 716

Academic Detailing Community Resource Teams, 678, 680–681

Accelerated junctional rhythm, electrocardiography in, 911

Acetabulum, labral tears of, 191–193

Acetaminophen, for pain, in end-of-life care, 816

Acetazolamide, for central sleep apnea, 370

Achilles tendon injuries, 120–121

Acromegaly, central sleep apnea in, 367

Actigraphy, wrist
in delayed sleep phase circadian sleep disorders, 455
in sleep deprivation, 483–484

Activities of daily living
assessment of, 661–663, 832–833. *See also* Incremental geriatric assessment.
description of, 620

Activity Counseling Trial, 952

Acute coronary syndrome, 1066–1073. *See also* Angina, unstable; Myocardial infarction.
assessment of, 1066–1067
definition of, 1066
diagnosis of, 932–934
management of, 1067–1073

Adductor muscle, strains of, 166–168

Adenoid hypertrophy, obstructive sleep apnea in, 557

Adenosine, for stress testing, in coronary artery disease, 936, 1063

Adjustment disorder with depressed mood, 784

Adolescents, sleep disorders in, 557, 559–560

Adrenergic blockers, sleepiness due to, 502

Adult Treatment Panel, coronary artery disease risk estimation guidelines of, 996, 1001–1002

Advance directives, 813
in dementia, 674–675
shared decision making in, 647–648

Advanced sleep phase circadian rhythm sleep disorders, 456–457, 577–579

AFFIRM (Atrial Fibrillation Follow-up Investigation of Rhythm Management), 1097–1098

Age-related maculopathy, in hypertension, 729

Aging
coronary artery disease risk and, 963–964
demographics of, 600–601, 795–796
nutritional changes with, 659–660
versus dementia, 611

Air pollution, coronary artery disease risk in, 972

Airway obstruction, apnea in. *See* Obstructive sleep apnea-hypopnea syndrome.

Akathisia, versus restless legs syndrome, 440

Alcohol ingestion and abuse
cardiovascular disease and, 966
depression in, 784
malnutrition in, 664
screening for, 633, 849–850
sedation in, 495
sleep disorders in, 324–325, 547

0095-4543/05/$ - see front matter © 2005 Elsevier Inc. All rights reserved.
doi:10.1016/S0095-4543(05)00097-7

ALCOVE-2 (Assessing Care of Vulnerable Elderly Adults in the office setting), 602–605

Aldosterone antagonists
for congestive heart failure, 1124
for hypertension, 737

Allen's test
in hand injuries, 81, 94–95
in wrist injuries, 27

ALLHAT (Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial), 1019–1020

Alprazolam, potentially inappropriate prescription of, 765

Alteplase, for coronary artery disease, 1071

Alzheimer's Association, resources of, 675–676

Alzheimer's disease
awareness of, 614
care of, **671–682**
barriers to, 672–674
community resources for, 675–678
end-of-life, 678
interactions throughout, 674–678
optimal, 671–672
physician role in, 671–674
services for, new strategies for, 678–681
caregivers in, 674–675, 680–681
demographics of, 600–601
diagnosis of, 608–611, 614, 674, 838
differential diagnosis of, 616
early interventions for, 611
empowerment in, 615
epidemiology of, 607–608
health maintenance in, 605–607
office demands and, 600–605
reimbursement issues in, 613–614
safety in, 616
sleep disorders in, 536–537
treatment of, 615–616
versus normal aging, 611
warning signs of, 612–613

Alzheimer's Disease Education and Referral (ADEAR) Association, 676

Alzheimer's Service Coordination Program, 678–680

American Academy of Sleep Disorders, insomnia nosology of, 377–378

American College of Cardiology
acute chest pain guidelines of, 932–934
atrial fibrillation classification of, 1085
exercise electrocardiography guidelines of, 934–935
functional classification of, 1121

American Heart Association
acute chest pain guidelines of, 932–934
atrial fibrillation classification of, 1085
exercise electrocardiography guidelines of, 934–935
functional classification of, 1121
Multiple Risk Factor Assessment Equation of, 1019

Amiloride, for hypertension, 739–740

Amino acids, as ergogenic aids, 287

Amiodarone
for atrial fibrillation, 1094–1095, 1100–1101
potentially inappropriate prescription of, 767

Amitriptyline
for depression, 788
for insomnia, 383, 498
potentially inappropriate prescription of, 764

Amlodipine
for coronary artery disease, 1064
for hypertension, 742–743, 1020

Amoxapine, for insomnia, 499

Amphetamines
for narcolepsy, 397–399, 500
potentially inappropriate prescription of, 767–768

Amyotrophic lateral sclerosis, sleep disorders in, 540

Anabolic steroids, as ergogenic aids, 280–282

Anatomic snuffbox, examination of, 26

Androstenedione, as ergogenic aid, 282

Angina
atypical, 1057–1058
grading of, 1059
hypertension in, 1021–1022
Prinzmetal's, 1064
stable
computed tomography in, 1063
echocardiography in, 1060
electrocardiography in, 1059–1060
stress testing in, 1060–1063
treatment of, 1063–1070, 1074
versus unstable angina, 1058–1059
typical, 1057

unstable
mortality prediction in, 1062
risk assessment in, 1067
treatment of, 1067–1070, 1074
versus stable angina, 1058–1059

Angiography, coronary
for coronary artery disease, 1065
in congestive heart failure, 1120

Angiotensin II, in congestive heart failure, 1116–1117

Angiotensin II receptor blockers
for congestive heart failure, 1123
for hypertension, 1021

Angiotensin-converting enzyme inhibitors
for atrial fibrillation, 1102
for congestive heart failure, 1122–1123
for coronary artery disease, 1065, 1069, 1073
for hypertension, 737, 739–741, 1020–1023

Ankle injuries
Achilles tendon, examination in, 120–121
anatomic considerations in, 105–106
chronic pain after, 143
examination in, 116–122
lateral instability, 142
sprains, 133–143
anatomic considerations in, 134–135
epidemiology of, 133
examination in, 117–119, 135–136
history in, 135
lateral, 136–137
medial, 142–143
osteochondritis dissecans in, 270, 272
prevention of, 141
radiography in, 137
rehabilitation in, 138–141
syndesmosis, 142–143
treatment of, 137–141

Anorexia nervosa, sleep disorders in, 546–547

Antalgic gait, in athletes, 109

Anterior drawer test, in ankle sprains, 118, 136–137

Antiarrhythmic agents, for atrial fibrillation, 1094–1095

Anticholinergic drugs
for urinary incontinence, 716
potentially inappropriate prescription of, 766

Anticoagulants
for atrial fibrillation, 1093, 1095–1096, 1099–1100
for coronary artery disease, 1067–1068

Anticonvulsants, sleepiness due to, 503

Antidepressants
for depression, 786–790
for insomnia, 381–383, 385–386, 498–499
for narcolepsy, 399–400
sleepiness due to, 503

Antidiarrheal agents, sleepiness due to, 504

Antidromic atrioventricular re-entry tachycardia, electrocardiography in, 906–907

Antiemetics, sleepiness due to, 504

Antihistamines
for insomnia, 382–383, 495, 497
potentially inappropriate prescription of, 766
sleepiness due to, 502

Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT), 737, 1019–1020

Antimuscarinic drugs, sleepiness due to, 502

Antiplatelet agents, for coronary artery disease, 1067–1068

Antipsychotics, sleepiness due to, 504

Antispasmodic drugs, sleepiness due to, 502

Antithrombotic agents, for coronary artery disease, 1067–1068

Antitussives, sleepiness due to, 504

Anxiety
in end-of-life care, 820–821
screening for, 633
sleep disorders in, 377, 544–546

Anxiolytics, sleepiness due to, 504

Aortic valve
regurgitation of, 1110, 1112
stenosis of, 1109–1112

Apheresis, low-density lipoprotein, 1050–1051

Apnea, sleep. *See* Central sleep apnea; Obstructive sleep apnea-hypopnea syndrome; Sleep apnea.

Apnea-hypopnea index, 330–331
heart disease and, 513
in obstructive sleep apnea, 344
versus body mass index, 332

Apolipoprotein(s), 1028

Apolipoprotein B, excess of, 1041

Apolipoprotein B_{100}, defective, 1005, 1038

Apophysitis
foot, 148–150
hip, in pediatric patients, 173–174

Appetite, loss of, 663–664

Arcade of Frohse, radial nerve compression at, 5–6

Arches, of foot, examination of, 108

Area Agencies on Aging, resources of, 676

Arginine, as ergogenic aid, 287

Arousal(s), from sleep, confusional, 306, 314–315, 416

Arousal index, in obstructive sleep apnea, 344

Arrhythmias, cardiac
electrocardiography for. *See* Electrocardiography; *specific arrhythmias.*
in congestive heart failure, 1119
in obstructive sleep apnea, 335
in rapid eye movement sleep–related sinus arrest, 420–421

Arthroscopy
in osteochondral injuries, 263–264, 266, 272
in triangular fibrocartilage complex injury, 45

Articular cartilage, injuries of. *See* Osteochondral injuries.

Aspiration, of wrist ganglion, 63

Aspirin
for atrial fibrillation, 1099–1100
for coronary artery disease, 1064, 1067–1068, 1073

Assessing Care of Vulnerable Elderly Adults in the office setting (ALCOVE-2), 602–605

Assisted living facilities, 796

Assistive devices, for fall prevention, 695–696

Asthma
gastroesophageal reflux with, 525–526
sleep disorders in, 518–521, 555

Asystole, in rapid eye movement sleep–related sinus arrest, 420–421

Atenolol
for coronary artery disease, 1064, 1068
for hypertension, 739–740

Atherosclerosis
blood pressure measurement in, 731–732
coronary. *See* Coronary artery disease.
coronary artery disease risk in, 966
erectile dysfunction in, 980–984
high-density lipoprotein prevention of, 1030
in hyperlipidemia, 1027–1028, 1030–1031, 1038–1039
in hypertension, 729

Atlantoaxial instability, athletic participation and, 239–240

Atrial appendage, isolation of, for atrial fibrillation, 1102

Atrial fibrillation, **1083–1107**
causes of, 1084
classification of, 1085
definition of, 1083
electrocardiography in, 905–908
electrophysiological mechanism of, 1085–1086
epidemiology of, 1083–1085
first detected
diagnosis of, 1087
indications for hospitalization in, 1088
initial workup of, 1088–1089
natural history of, 1088
treatment of, 1089–1096
lone, 1100
pathophysiology of, 1085–1086
postoperative, 1096
prognosis for, 1086–1087
recurrent
rate versus rhythm control in, 1096–1098
treatment of, 1096–1103
referral in, 1103
triggers of, 1084

Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM), 1097–1098

Atrial flutter, electrocardiography in, 908–909

Atrial kick, loss of, in atrial fibrillation, 1086

Atrial tachycardia, electrocardiography in, 907

Atrioventricular block, electrocardiography in, 911–915

Atrioventricular nodal re-entry tachycardia, electrocardiography in, 906

Atrioventricular node, ablation of, for atrial fibrillation, 1102

Atrioventricular re-entry tachycardia, electrocardiography in, 906–907

Atrium, enlargement of, electrocardiography in, 903–904

Attention-deficit hyperactivity disorder, sleep disorders with, 555

Australian National Blood Pressure Study, 740, 1020

Autologous chondrocyte implantation, for knee osteochondral injuries, 267–268

Automatic behavior, in narcolepsy, 390

Autonomic dysfunction, in sleep deprivation, 479

Autonomy, shared decision making and, 645–646, 648–649

Avascular necrosis
of hip, 194–196
of lunate, 55–56

Avulsion fractures, of hip, in pediatric patients, 173–174

B

Back pain, in athletes, **201–229**
causes of, 203, 215
diagnosis of, 202
equipment causing, 202
in disc injuries, 214–218
in muscular strain, 213–216
in sacral stress fracture, 220
in sacroiliitis, 220
in Scheuermann's disease, 203, 210
in scoliosis, 211–212
in spinal stenosis, 215, 219–221
in spondylolisthesis, 203, 208–210
in spondylolysis, 204–208
in vertebral fractures, 203, 212–213, 218–219
mature, 214–220
prevalence of, 201, 214–215
risk factors for, 201–202
sports related to, 203, 215
symptoms of, 215
treatment of, 220–225
young, 202–214

Balance
assessment of, for fall risk, 688–690
exercises for, for fall prevention, 693–695

Bangungut (sudden unexplained nocturnal death syndrome), 428–429

Barbiturates
for insomnia, 497
potentially inappropriate prescription of, 766
sleepiness due to, 503–504

BEARS mnemonic, for sleep disorder evaluation, 319

Bedwetting, 505, 554–555

Beers criteria, for drug prescription, in elderly persons, 762–768

Behavioral disorders, sleep disorders with, 324

Behavioral modification
counseling for, 5A's for, 950, 957
for insomnia, 568–569
for urinary incontinence, 709–714

Belladonna, potentially inappropriate prescription of, 766

Bennett fracture, 102

Benzodiazepines
for end-of-life care, 818, 823–824
for insomnia, 383–384, 494, 496
for parasomnias, 554
for restless legs syndrome, 444
potentially inappropriate prescription of, 765
sleepiness due to, 503–504

Bereavement
depression in, 782
in end-of-life care, 824–825

Berndt and Harty classification, of talar osteochondral lesions, 153

Beta-blockers
as ergogenic aids, 286–287
for atrial fibrillation, 1091–1093, 1098
for congestive heart failure, 1123–1124
for coronary artery disease, 1064, 1068–1069, 1073
for hypertension, 737, 1022
sleepiness due to, 502

Biatrial enlargement, electrocardiography in, 904

Biceps muscle, distal, injury of, 12

Bicuspid aortic valves, 1109

Bifascicular block, electrocardiography in, 920

Bile acid sequestrants, for hyperlipidemia, 1048

Biomechanics, of sports injuries, **245–251**
anatomical variants and, 248
jumping, 247–249
kicking, 249–251
running, 245–247

Bipolar disorder, sleep disorders in, 386, 543–544

Bisoprolol, for coronary artery disease, 1064

Bivalirudin, for coronary artery disease, 1068

Blindness, nonentrained circadian rhythm sleep disorders in, 458–459

Blood doping, for sports performance, 282–283

Blood pressure
age-related changes in, 727–728
circadian variation of, 731
high. *See* Hypertension.
measurement of, 731–732
ambulatory, 734
self-recorded, 735
monitoring of, ambulatory, 1014–1016
orthostatic changes in, fall risk in, 689
postprandial, 732–733

Blood Pressure lowering Treatment Trialists' Collaboration, 1020

Body alignment, observation of, in foot and ankle injuries, 107–108

Body mass index, versus apnea-hypopnea index, 332

Bone scan
in osteitis pubis, 187
in osteochondral injuries, 261
in scaphoid fracture, 50–51
in wrist injuries, 31–32

Botulinum toxin, for bruxism, 423

Boutonnière finger, 86, 99

Boxer's knuckle, 100

Brachial artery
anatomy of, 3
ultrasonography of, in endothelial function evaluation, 941–942

Brachial plexus injuries, in athletics, 234

Bracing
for back pain, 224
for osteochondral injuries, 264

Bradyarrhythmias, in sick sinus syndrome, 905

Brain natriuretic peptide, coronary artery disease and, 973

Breathing, sleep-disordered, screening for, 315

Breathlessness, in end-of-life care, 818

Broad band dyslipoproteinemia, 1041

Brugada criteria, for ventricular tachycardia, 922

Bruxism, 314, 421–424

B-type natriuretic peptide, in congestive heart failure, 1119

Bulimia, sleep disorders in, 546–547

Bundle branch blocks, electrocardiography in, 916–919

Bunion, 128

Bunionette, 128

Bupivacaine, for DeQuervain's tenosynovitis, 95

Bupropion
for depression, 787–788
for insomnia, 499
for smoking cessation, 949

Bursitis, iliopsoas, 169–171

Buspirone, for insomnia, 499

C

Cabergoline, for restless legs syndrome, 443

Caffeine
as ergogenic aid, 279–280
for sleep deprivation, 486
insomnia due to, 567–568

CAGE questionnaire, for alcoholism screening, 850

Calcaneonavicular coalition, 151–152

Calcaneus, apophysitis of (Sever's disease), 126, 149–150

Calcium, in coronary arteries, computed tomography in, 938–939, 941, 971

Calcium channel blockers
for atrial fibrillation, 1091–1093, 1098
for coronary artery disease, 1064–1065, 1068

Canadian Cardiovascular Society, angina classification of, 1058–1059

Canadian C-Spine Rule, for cervical injury evaluation, 232–233

Cane, for fall prevention, 695–696

Capacity, for decision making
determination of, 649–650
lack of, 651–652

Capitate
anatomy of, 21
fracture of, 58

Capitellum, osteochondral injuries of, 273

Captopril, for congestive heart failure, 1123

Carbamazepine, for nocturnal paroxysmal dystonia, 417–418

Carbidopa-levodopa, for restless legs syndrome, 441–443, 576

Cardiac arrhythmias. *See also* Electrocardiography; *specific arrhythmias.*
in congestive heart failure, 1119
in obstructive sleep apnea, 335
rapid eye movement sleep–related, 420–421

Cardiac rehabilitation, in coronary artery disease, 1075

Cardiomyopathy
hypertrophic, murmurs in, 1110
ischemic, heart failure in, 1116

Cardiopulmonary resuscitation, in cervical spinal injuries, 231–233

Cardiorespiratory processes, in sleep, 323–324

Cardiovascular disease. *See also* Coronary artery disease; Myocardial infarction; *specific diseases, eg,* Congestive heart failure.
depression in, 782
electrocardiography in, **901–930**
erectile dysfunction in, **777–993**
in hyperlipidemia, **1027–1055**
in hypertension, 728–731, **1011–1025**
obstructive sleep apnea and, 333–335
prevention of
community interventions for, **865–881**
lifestyle interventions for, **947–961**
office systems for, **883–900**
risk factors for, **963–976**
sleep and, 511–518
valvular, **1109–1114**. *See also specific disorders.*

Cardioversion, for atrial fibrillation, 1093–1095

Caregiver(s)
burden of, assessment of, 851–853
in dementia, 674–675, 680–681

Caregiver strain questionnaire, 853

Carotid artery, injury of, in sports, 240–241

Carpal bones, anatomy of, 18–21

Carpal tunnel, anatomy of, 21

Carpal tunnel syndrome, 64–65

Carpometacarpal joint, anatomy of, 72

Cartilage
composition and function of, 254–255
injuries of. *See* Osteochondral injuries.

Carvedilol, for congestive heart failure, 1123–1124

Cast
for distal radial fracture, 46, 48
for hamate fracture, 56
for scaphoid fracture, 50–53

Cataplexy, 558
narcolepsy and, 390, 394, 399–402
treatment of, side effects of, 506

Catastrophic accidents, due to sleep deprivation, 481–482

CATCH (Child and Adolescent Trial for Cardiovascular Health), 869, 873–874

Catheter ablation, for atrial fibrillation, 1103

Central hypoventilation syndrome, 552–553

Central sleep apnea, **361–374**
age factors in, 365
causes of, 362
classification of, 361–362
clinical features of, 367–368
definition of, 361
development of, 363–367
diagnosis of, 368
gender differences in, 365
hypercapnic, 361–362, 368–369
idiopathic, 367
in congestive heart failure, 366–367
in metabolic disorders, 367
in stroke, 367
initiation of, 363–364
nonhypercapnic, 362, 368–369
overview of, 308
pathophysiology of, 361–362
perpetuation of, 364–365
sleep state and, 365–366
treatment of, 368–371
vulnerability to, 365–367

Certification, for electrocardiogram interpretation, 926–927

Cervical injuries, **231–243**
differential diagnosis of, 241
discogenic, 238–239
emergency, in athletic event, 231–233
functional stenosis and, 235–236
in Down syndrome, 239–240
in rheumatoid arthritis, 240
in spear tackling, 233
return-to-sport recommendations in, 236
spinal cord neuropraxia, 234–236
spinal stenosis and, 235
stingers, 234
strains and sprains, 237–238
vascular, 240–241

CHAD (Community Syndrome of Hypertension, Atherosclerosis and Diabetes) program, 866–867

Charley horse, 417, 440

Chest pain
acute, diagnosis of, 932–934
anginal. *See* Angina.
atypical, 932
cardiac etiologies of, 1057
evaluation of, 1057–1058
nonanginal, 932
noncardiac causes of, 1058

Chest radiography, in atrial fibrillation, 1088–1089

Cheyne-Stokes respiration, in congestive heart failure, 515–518

Child and Adolescent Trial for Cardiovascular Health (CATCH), 869, 873–874

Chloral hydrate, for insomnia, 493, 497

Chlorpheniramine, potentially inappropriate prescription of, 766

Chlorpropamide, potentially inappropriate prescription of, 766

Chlorthalidone, for hypertension, 738, 1020

Cholestelam, for hyperlipidemia, 1048

Cholesterol
disorders of. *See* Hyperlipidemia.
excess of, in familial hypercholesterolemia, 1004–1005
measurement of, 1033–1034

Cholesterol absorption inhibitors, for hyperlipidemia, 1047–1048

Cholesterol efflux regulatory protein, 1030

Cholesterol ester transfer protein, 1030
inhibitors of, for hyperlipidemia, 1050

Cholestyramine, for hyperlipidemia, 1048

Chondral lesions, definition of, 253

Chondrocytes, implantation of, for knee osteochondral injuries, 267–268, 270

Chronic Care Network for Alzheimer's Disease, 678–680

Chronic illness, depression in, 779–781

Chronic obstructive pulmonary disease, sleep disorders in, 521–524

Chronotherapy
for advanced phase circadian sleep disorders, 457
for delayed sleep phase circadian sleep disorders, 455, 558
for irregular sleep–wake disorders, 459

Chylomicronemia syndrome, 1040

Chylomicrons and chylomicron remnants, 1028–1029

Cimetidine, potentially inappropriate prescription of, 768

Circadian rhythm, development of, 551

Circadian rhythm sleep disorders, 314–315, **449–473**
advanced sleep phase type, 456–457, 577–579
delayed sleep phase type, 453–456, 557–558
irregular sleep–wake type, 459
jet lag type, 460–462
nonentrained type, 458–459
physiologic basis for, 320–321, 449–452
shift work type, 462–465
treatment of
safety issues in, 465–466
side effects of, 506

Citalopram
for depression, 787
for insomnia, 499

Claw toe, 129

Clidinium, potentially inappropriate prescription of, 766

Clinical trials, of drugs, for elderly persons, 760–762

Clock drawing test, 627, 834

Clonazepam
for insomnia, 496

for rapid eye movement sleep behavior disorder, 420, 582–583
for restless legs syndrome, 444

Clonidine
for restless legs syndrome, 576
potentially inappropriate prescription of, 768

Clopidogrel, for coronary artery disease, 1064, 1068, 1073

Clorazepate, for insomnia, 494

Closed reduction, of distal radial fracture, 46, 48

Coach's finger, 71, 101–102

Cockroft-Gault equation, for creatinine clearance, in elderly persons, 756–757

Codeine, for restless legs syndrome, 576

Coeur en Santé Saint Henri Project, 873

Cognitive function
antihypertensive therapy effects on, 743
assessment of, 626–627, 834–840
hypertension effects on, 729–731

Cognitive-behavioral therapy, for insomnia, 382

Colestipol, for hyperlipidemia, 1048

Collagen vascular disease, cardiovascular disease in, 971

Collateral ligaments, at elbow
anatomy of, 2
injury of, 8–10

Colles' fracture, 46, 48

Communication
in end-of-life care, 813–814
in nursing facility setting, 802–803

Community interventions, for cardiovascular disease, **865–881**
components of, 877–878
historical perspective of, 866–875
modest impact of, 875–877

Community resources, for dementia care, 675–681

Community Syndrome of Hypertension, Atherosclerosis and Diabetes (CHAD) program, 866–867

Community-oriented primary care model, 874–875

Compartment syndrome, of lower extremity, 115–116

Complementary medicine, drug prescription precautions with, 769–770

Complete (third-degree) heart block, electrocardiography in, 914–915

Compliance, with drug treatment, barriers to, in elderly persons, 763, 768–769

Compression fractures, vertebral, in athletes, 215, 218–219

Computed tomography
in coronary artery disease, 937–939, 941, 964, 1063
in foot fractures, 146
in osteochondral injuries, 261
in scaphoid fracture, 51
in spondylolysis, 206
in tarsal coalition, 152
in triangular fibrocartilage complex injury, 45
in ulnar collateral ligament injury, 9
in wrist injuries, 32

Confusional arousals, from sleep, 314–315, 416

Congestive heart failure, **1115–1129**
apnea-hypopnea index in, 513
central sleep apnea in, 366–367, 371
Cheyne-Stokes respiration in, 515–518
clinical likelihood of, 1118–1119
diagnosis of, 1117–1121
diastolic, 1117, 1125
economic impact of, 1115–1116
epidemiology of, 1115–1116
hypertension with, 729, 1022
natural history of, 1116–1117
obstructive sleep apnea in, 335
pathophysiology of, 1116–1117
referral in, 1125–1126
spectrum of, 1117
staging of, 1121
treatment of, 1121–1127
collaborative, 1125–1126
in diastolic dysfunction, 1125
nonpharmacologic, 1121–1122
palliative care in, 1126–1127
pharmacologic, 1122–1125
Vesper's curse in, 440

Constipation, in end-of-life care, 819–820

Continuous positive airway pressure
for asthma, 520–521
for central sleep apnea, 369, 371
for Cheyne-Stokes respiration, 517–518
for obstructive sleep apnea, 346–349, 513–514, 573

COPD (chronic obstructive pulmonary disease), sleep disorders in, 521–524

CORIS (Coronary Risk Factor Study), 867

Cornell scale for depression in dementia, 847–848

Coronary angiography
in congestive heart failure, 1120
in coronary artery disease, 1065–1066

Coronary artery bypass graft surgery, 1065–1066, 1069–1070

Coronary artery disease, **931–946**
acute coronary syndrome in. *See* Acute coronary syndrome; Angina; Myocardial infarction.
cardiac rehabilitation in, 1075
diagnosis of
acute chest pain and acute coronary syndrome and, 932–934
computed tomography in, 937–939
exercise electrocardiography in, 934–935
nuclear perfusion imaging in, 935–936
pretest probability and, 932
stress echocardiography in, 936–937, 1061–1063
endothelial dysfunction in, 980–981
erectile dysfunction in, 982–986
family history of, **995–1010**
dyslipidemias, 1004–1005. *See also* Hyperlipidemia.
importance of, 996–997
patient identification and, 997
patient reactions to, 997, 1000–1001
preventive treatment in, 1005–1006
relative risk of, 964, 997–1000
risk estimation in, 1001–1004
hypertension causing, 1011, 1019
myocardial infarction in. *See* Myocardial infarction.
obstructive sleep apnea and, 335
premature, 997, 1004
prevention of, **947–961**
counseling on, 957
diet in, 953–955
exercise in, 950–953
family history factors and. *See* Coronary artery disease, family history of.
multiple behavior interventions in, 955–957
office systems for, **883–900**
screening in, 939–942
smoking cessation in, 947–950
risk factors for, 939–940, **963–976**, 1034–1037
age, 963–964
air pollution, 972
alcohol ingestion, 966
brain natriuretic peptide, 973
calcification, 971
collagen vascular disease, 971
C-reactive protein, 968
diabetes mellitus, 965
dietary, 953–955, 966, 1044
endothelial dysfunction, 941–942, 971–972, 980–984
family history, **995–1010**
fibrinogen, 970–971
gender, 964
heart rates, 971
homocysteine, 968–969
hyperlipidemia. *See* Hyperlipidemia.
hypertension, 949, 965
hyperuricemia, 972
infections, 972
iron overload, 973
kidney disease, 967
lipid abnormalities, 965
lipoprotein(a), 969–970
metabolic syndrome, 967
obesity, 966
periodontal disease, 972
physical inactivity, 965–966
psychosocial, 967
small for gestational age, 973
smoking, 947–950, 964, 1073
socioeconomic, 973
ventricular hypertrophy, 966
screening for, 884–885
coronary artery calcium measurement in, 941
endothelial function measurement in, 941–942
exercise electrocardiography in, 940
risk prediction in, 939–940
treatment of
follow-up in, 1073–1076
in myocardial infarction, 1067–1071
in post-acute phase, 1071–1073
in unstable angina, 1067–1070
lifestyle modifications in, 1073–1074
pharmacologic, 1063–1065
principles of, 1063
revascularization in, 1065–1066

Coronary Risk Factor Study (CORIS), 867

Corticosteroids
for carpal tunnel syndrome, 64
for nocturnal asthma, 521
for pain, in end-of-life care, 817
hip avascular necrosis due to, 195
injection of
for DeQuervain's tenosynovitis, 38, 95
for extensor carpi ulnaris tendonopathy, 40
for iliopsoas strains and bursitis, 170–171
for lateral epicondylitis, 5
for osteitis pubis, 188

Cough, in sleep-related abnormal swallowing syndrome, 425

Counseling, for behavioral change, 5A's for, 950, 957

Cramps, muscle, nocturnal, 417, 440

C-reactive protein, elevated
coronary artery disease risk in, 968
in sleep deprivation, 479

Creatine, as ergogenic aid, 283–285

Creatinine clearance, in elderly persons, 756–757

Cubital tunnel syndrome, 10–11

Cyclandelate, potentially inappropriate prescription of, 766, 768

Cyclist's palsy, 65

Cyproheptadine, potentially inappropriate prescription of, 766

Cyst, ganglion, of wrist, 63

Cytokines, in obstructive sleep apnea, 512

D

Darvocet, for restless legs syndrome, 576

Day care, adult, in dementia, 677

Death and dying
active process of, 823–824
family care after, 824
from adverse drug reactions, 758
from falls, 683–685
grief and bereavement after, 782, 824–825
in sudden infant death syndrome, 552
in sudden unexplained nocturnal death syndrome, 428–429

Debridement, for knee osteochondral injuries, 266

Decision making
capacity for, evaluation of, 633–634
in end-of-life care, 813
shared. *See* Shared decision making.

Defibrillators, implantable, for coronary artery disease, 1075–1076

Dehydration, in end-of-life care, 819

Dehydroepiandrosterone, as ergogenic aid, 282

Delayed sleep phase circadian rhythm sleep disorders, 453–456, 557–558

Delirium
diagnostic criteria for, 840
in end-of-life care, 823–824

Dementia, **599–618**
awareness of, 614
care of, **671–682**
barriers to, 672–674
community resources for, 675–681
end-of-life, 678
interactions throughout, 674–678
optimal, 671–672
physician role in, 671–674
services for, new strategies for, 678–681
caregivers in, 674–675, 680–681
demographics of, 600–601
depression with, assessment of, 633, 847–848
diagnosis of, 608–611, 614, 674, 838–839
differential diagnosis of, 616
early interventions for, 611
empowerment in, 615
epidemiology of, 607–608
family fears and stress in, 614
health maintenance in, 605–607
hypertension and, 730
office demands and, 600–605
pessimism about, 613
reimbursement issues in, 613–614
reversible, 616
safety in, 616
sleep disorders in, 536–539
treatment of, 615–616
versus normal aging, 611
warning signs of, 612–613

Dental appliances, for bruxism, 423

Department of Veterans Affairs Cooperative Study, of hypertension, 742

Depression, **777–792**
criteria for, 633, 784
dementia with, assessment of, 633, 847–848
diagnosis of, 783–785, 845–846
in bereavement, 782
in end-of-life care, 821
prevalence of, 778–779
recognition of, 783
risk factors for, 779–782
screening for, 628–629, 632–633, 783, 844–848
sleep deprivation and, 479
sleep disorders in, 377, 385–386, 543
treatment of, 785–790

Deprivation, sleep. *See* Sleep deprivation.

DeQuervain's tenosynovitis, 37–38, 95

Desipramine
for depression, 787–788
for insomnia, 499

DETERMINE Warning Signs, for nutritional assessment, 661

Determine Your Nutritional Health Checklist, 661, 668

Developmental changes, in sleep physiology, 322–323

Dextroamphetamine, for narcolepsy, 500

Diabetes mellitus
coronary artery disease risk in, 965
dyslipidemia in, 1051–1052
erectile dysfunction in, 988
hypertension with, 741–742, 744, 1021

Diagnostic and Statistical Manual, 4th Edition, insomnia categories in, 378

Diamond-Forrester method, for coronary artery disease prediction, 932

DIAPPERS acronym, for urinary incontinence, 702–703

Diary
sleep, 483
urinary incontinence, 705, 719

Diazepam, for insomnia, 494

Dicyclomine, potentially inappropriate prescription of, 766

Diet. *See also* Nutrition.
for coronary artery disease prevention, 953–955, 966
for hyperlipidemia, 1044
hypertension and, 726
urinary incontinence and, 705

Dietary Approaches to Stop Hypertension (DASH) diet, 1016–1017

Dietary supplements, for hyperlipidemia, 1044–1045

Dietary Supplements Health and Education Act of 1994, ergogenic aids and, 277

Digoxin
for atrial fibrillation, 1091–1092
for congestive heart failure, 1125
potentially inappropriate prescription of, 765

Diltiazem
for atrial fibrillation, 1091–1093
for coronary artery disease, 1069
for hypertension, 739

Dim light melatonin onset, 451

Diphenhydramine
for insomnia, 495, 497
potentially inappropriate prescription of, 766

Dipyridamole
for nuclear perfusion imaging, in coronary artery disease, 936
for stress testing, in coronary artery disease, 1063
potentially inappropriate prescription of, 765

Disc(s), intervertebral, disorders of
cervical, 238–239
lumbar, 215–218

Discogenic neck pain, 238–239

Dislocation
of distal radioulnar joint, 42–44
of interphalangeal joints, 101–102

Disopyramide
for atrial fibrillation, 1095
potentially inappropriate prescription of, 765

Distal posterior interosseous nerve syndrome, 65

Diuretics
for congestive heart failure, 1124–1125
for hypertension, 737, 741–742

Dobutamine
for nuclear perfusion imaging, in coronary artery disease, 936
for stress echocardiography, in coronary artery disease, 936–937

Documentation, of nursing facility care, 807–808

Dofetilide
for atrial fibrillation, 1101
potentially inappropriate prescription of, 1095

Do-not-resuscitate status, 813

Dopamine agonists
for restless legs syndrome, 442–444
sleepiness due to, 502

Doping, for sports performance. *See* Ergogenic aids.

Doppler studies, of blood pressure, in extremities, 733

Dorsal intercalated segment instability, 62

Down syndrome, athletic participation in, neck considerations in, 239–240

Doxazosin
for hypertension, 1020
potentially inappropriate prescription of, 768

Doxepin
for depression, 787
for insomnia, 499
potentially inappropriate prescription of, 765

Drawer test, anterior, in ankle sprains, 118, 136–137

Dreams, frightening (nightmares), 314–315, 416–417, 554

Driving, accidents during, due to sleep deprivation, 333, 481, 484–486, 559–560

Drug(s)
adverse reactions to, in elderly persons, 757–760
anorexia due to, 663–664
arrhythmias due to, 924
depression due to, 781
falls due to, 690
hypertension due to, 726–727
interactions of, in elderly persons, 757–760
prescribing, for elderly persons, **755–775**
adverse drug reactions in, 757–760
Beers criteria for, 762–768
challenges in, 755
complementary medicine and, 769–770
compliance barriers in, 763, 768–769
cost considerations in, 768–769
drug interactions in, 757–760, 770
drug review in, 631, 771–772
drug trials for, 760–762
errors in, 758–760
in nursing facility setting, 803
multiple, 759
over-the counter drug use and, 769–770
pharmacodynamics and pharmacokinetics of, 755–757
practical strategies for, 771–773
SEA-squared model for, 772–773
undertreatment with, 762
weaning in, 771–772
sleep effects of, **491–509**
daytime sleepiness as, 502–504
disordered sleep as, 492
for alertness induction, 500–501
for cataplexy, 506
for circadian rhythm sleep disorders, 506
for daytime sleepiness, 498–500
for enuresis, 505
for insomnia, 492–498
for parasomnias, 501, 505
for periodic limb movement disorder, 505–506
for restless legs syndrome, 505–506
respiratory, 505
urinary incontinence due to, 706

Drunkenness, sleep, 402–404

Dying. *See* Death and dying.

Dynamic therapy, for depression, 786

Dysbetalipoproteinemia, familial, 1005

Dyslipidemia. *See* Hyperlipidemia.

Dyslipoproteinemia, broad band, 1041

Dyspnea
in congestive heart failure, 1117–1118
in end-of-life care, 818

Dysthymia, definition of, 784

Dystonia, nocturnal paroxysmal, 417–418

E

Eating disorders, sleep disorders in, 546–547

Echocardiography
in aortic regurgitation, 1112
in atrial fibrillation, 1088–1089
in congestive heart failure, 1119–1120
in mitral regurgitation, 1113
in mitral stenosis, 1113

Echocardiography (*continued*)
in stable angina, 1060
stress, in coronary artery disease, 936–937, 1061–1063

Economic issues, in sleep disorders, 305–306, 377

Edema, in congestive heart failure, 1118, 1122

Elbow injuries, **1–16**
anatomic considerations in, 1–3
cubital tunnel syndrome, 10–11
distal biceps, 12
lateral epicondylitis (tennis elbow), 4–5
medial epicondylitis (golfer's elbow), 7–8
osteochondral, 273
posterior impingement, 13–15
posterior interosseous nerve syndrome, 7
posterolateral rotatory instability, 11
pronator syndrome, 3–4
radial tunnel syndrome, 5–6
triceps tendon, 13
ulnar collateral ligament, 8–10

Elderly persons
common diagnoses in, 601
dementia in. *See* Dementia.
depression in. *See* Depression.
drug prescribing for, **755–775**
end-of-life care for, 678, **811–828**
falls in, **683–697**
hyperlipidemia in, 1052–1053
hypertension in, **723–753**
in nursing facility setting, **793–810**
incremental assessment of, **619–643**
screening instruments for, **829–854**
shared decision making for, **645–658**
sleep disorders in, **563–586**
advanced sleep phase circadian rhythm disorders, 577–579
central sleep apnea, 365
history of, 565–566
insomnia, 566–569
obstructive sleep apnea, 569–574
rapid eye movement sleep behavior disorder, 579–583
restless legs syndrome, 574–577
versus normal sleep changes, 563–564
urinary incontinence in, **699–722**

Electrical cardioversion, for atrial fibrillation, 1094

Electrocardiography, **901–930**
baseline, 901–903
characteristics of
P wave, 901–910
PR interval, 910–915
Q wave, 922–923
QRS interval, 915–922
QT interval, 923
T wave, 923
U wave, 923–924
exercise, for coronary artery disease, 934–935, 940, 1060–1061
in atrial fibrillation, 905–908, 1087–1089
in atrial flutter, 908–909
in atrial tachycardia, 907
in atrioventricular block, 911–915
in atrioventricular nodal re-entry tachycardia, 906
in atrioventricular re-entry tachycardia, 906–907
in biatrial enlargement, 904
in fascicular block, 919–920
in hypercalcemia, 925
in hyperkalemia, 925
in hypertension, 734
in hypocalcemia, 925
in hypokalemia, 923, 925
in hypomagnesemia, 925
in junctional rhythm, 910–911
in left atrial enlargement, 903–904
in left bundle branch block, 918–919
in left ventricular hypertrophy, 915–916
in medication-induced arrhythmias, 924
in multifocal atrial tachycardia, 909–910
in myocardial infarction, 924, 926
in nonspecific intraventricular conduction delay, 921
in pre-excitation syndromes, 911
in premature atrial complex, 905
in premature ventricular contraction, 921
in right atrial enlargement, 903
in right bundle branch block, 916–918
in right ventricular hypertrophy, 916
in sick sinus syndrome, 905
in sinus arrhythmia, 905
in sinus tachycardia, 907
in stable angina, 1059–1060
in supraventricular tachycardia, 905–910
in ventricular tachycardia, 921–922
indications for, **901**–903
interpretation of, clinical competency in, 925–927
preoperative, 902–903

Electroencephalography
in sleep, 549–550
in sleep deprivation, 479–480

Electromyography
in obturator nerve entrapment, 175
in pronator syndrome, 4
in wrist injuries, 32

Electron beam computed tomography, in coronary artery disease, 937–939, 941, 964, 1063

Elson test, in hand injuries, 85–86

Emergency treatment, of cervical injuries, at athletic event, 231–233

Emotional problems, erectile dysfunction in, 988

Enalapril
for congestive heart failure, 1123
for hypertension, 738–739

End-of-life care, **811–828**
communication during, 813–814
coordination of, 814–815
goals of, 812–813
grief and bereavement in, 824–825
immediately after death, 824
in active dying process, 822–824
in congestive heart failure, 1126–1127
in dementia, 678
spiritual issues in, 821–822
symptom management in
physical, 815–820
psychosocial concerns in, 820–821

Endothelial dysfunction
erectile dysfunction in, cardiovascular disease risk and, 980–985
in coronary artery disease, 941–942, 971–972
in obstructive sleep apnea, 511–512

Endurance exercise, for fall prevention, 693

Enuresis, 505, 554–555

Environment, home, safety evaluation for, 636, 854

Epicondylitis
lateral (tennis elbow), 4–6
medial (golfer's elbow), 7–8

Epilepsy
nocturnal frontal lobe, 417–418
sleep disorders in, 541–542, 555

Eplerenone, for congestive heart failure, 1124

Eptifibatide, for coronary artery disease, 1068

Epworth Sleepiness Scale, 309–310
in elderly persons, 565–566
in narcolepsy, 401
in sleep deprivation, 483

Erectile dysfunction, cardiovascular disease and, **777–993**
as predicative factor, 982–984
associations, 978–979
endothelial dysfunction in, 980–982
evaluation of, 982–984
value of information acquired, 985–986
workup for, 985–987

Erections, painful, sleep-related, 425

Ergogenic aids, **277–292**
amino acids, 287
anabolic steroids, 280–282
beta blockers, 286–287
blood doping, 282–283
caffeine, 279–280
creatine, 283–285
erythropoietin, 282–283
growth hormone, 286
historical use of, 278
β-hydroxy-β-methylbutyrate, 287
identification of, 278–279
legislation on, 277
sodium bicarbonate, 287–288
stimulants, 279–280

Ergot mesyloids, potentially inappropriate prescription of, 766

Erythropoietin, as ergogenic aid, 282–283

Esmolol, for atrial fibrillation, 1091, 1093

Estazolam, for insomnia, 494, 496

Estrogens, potentially inappropriate prescription of, 768

Eszopiclone, for insomnia, 383, 494–496

Ethchlorvynol, for insomnia, 493, 497

Ethical issues
in decision making. *See* Shared decision making.
in nursing facilities, 803–804

European Working Party on High Blood Pressure in the Elderly, 738, 742

Exercise
for back pain, in athletes, 221–222
for congestive heart failure, 1122
for coronary artery disease prevention, 950–953, 1075
for fall prevention, 691–695
for hyperlipidemia, 1044
for hypertension, 1016–1017
for sleep deprivation, 485
for urinary incontinence, 709, 715–716

Exercise (*continued*)
in aortic stenosis, 1111
lack of, coronary artery disease risk in, 965–966

Exercise electrocardiography, in coronary artery disease, 934–935, 940, 1060–1061

Exercise nuclear perfusion imaging, in coronary artery disease, 936

Extensor carpi radialis muscle, anatomy of, 22–23

Extensor carpi ulnaris muscle, anatomy of, 22–23

Extensor carpi ulnaris tendon, injury of, 40–41

Extensor pollicis brevis tendon, De Quervain's tenosynovitis of, 37–38, 95

Extensor retinaculum, of wrist, anatomy of, 23

External rotation test, in ankle sprains, 119

Extracorporeal shock wave therapy, for plantar fasciitis, 156

Ezetimbe, for hyperlipidemia, 1047–1048

F

Falls, in elderly persons, **683–697**
definition of, 683
epidemiology of, 684–686
risk factors for, 604, 686–688
identification of, 634–635, 688–690
interventions for, 690–696

Familial combined hyperlipidemia, 1005, 1041

Familial defective apolipoprotein B_{100}, 1005, 1038

Familial dysbetalipoproteinemia, 1005

Familial high-density lipoprotein deficiency, 1042

Familial hypercholesterolemia, 1005, 1037–1038

Familial hypertriglyceridemia, 1005, 1040

Familial hypoalphalipoproteinemia, 1042

Familial lipoprotein lipase deficiency, 1005

Family
coronary artery disease history in. *See* Coronary artery disease, family history of.
of elderly person
care of, in active dying process, 822–824
communication with, in end-of-life care, 813–814
discussion with, on functional status, 635–636
with dementia, 614–615, 674–675, 680–681

Family Caregiver Alliance, 676

Fascicular block, electrocardiography in, 919–920

Fasting lipid profile, in hyperlipidemia, 1033

Fatigue
in congestive heart failure, 1117
in end-of-life care, 819

Fatigue fractures. *See* Stress fractures.

Felodipine
for coronary artery disease, 1064
for hypertension, 739

Felon, 78

Femoral cutaneous nerve, lateral, entrapment of, 178

Femoral nerve, entrapment of, 175–176

Femur
head of
avascular necrosis of, 194–195
cartilage lesions in, 193–194
neck of, stress fractures of, 189–191

Fenofibrate, for hyperlipidemia, 1049

Fentanyl, for pain, in end-of-life care, 817

Ferritin, deficiency of, restless legs syndrome in, 439, 441

Ferrous sulfate, potentially inappropriate prescription of, 766

Fetal tissue transplantation, for narcolepsy, 401

Fibrates, for hyperlipidemia, 1048–1049

Fibrillation, atrial. *See* Atrial fibrillation.

Fibrinogen, elevated, coronary artery disease risk in, 970–971

Finger extension test, in wrist injuries, 29–30

Finger injuries
anatomic considerations in, 71–73, 92–93
boutonnière finger, 86, 99
coach's finger, 71, 101–102

distal phalangeal, 102
fractures, 100–102
history of, 91–92
jersey finger, 84–85, 98–99
mallet finger, 85, 97–98
physical examination in, 75–88, 93–95
Elson test in, 85–86
Finkelstein test in, 87
flexor digitorum profundus and superficialis testing in, 84–85
inspection in, 75, 77
mallet finger test in, 85
neurovascular, 80–81, 94
palpation in, 77–78
range of motion assessment in, 78–80
strength testing in, 81–82, 84
volar plate assessment in, 86
pseudoboutonnière deformity, 86, 99
swan-neck deformity, 86
trigger finger, 99

Finkelstein's test
in DeQuervain's tenosynovitis, 95
in hand injuries, 87
in wrist injuries, 27, 38

First metatarsal rise test, in hindfoot injuries, 124

5A's, for behavior change counseling, 886–893, 950, 957

Flecainide, for atrial fibrillation, 1094–1095

Flexor carpi radialis muscle, anatomy of, 22–23

Flexor carpi radialis tendon, injury of, 42

Flexor carpi ulnaris muscle, anatomy of, 22

Flexor carpi ulnaris tendon, injury of, 42

Flexor digitorum profundus muscle, testing of, in injury, 84–85, 94

Flexor digitorum superficialis muscle, testing of, in injury, 84–85, 94

Flexor hallucis longus tendonitis, 125

Flexor retinaculum, of wrist, anatomy of, 23

Fluoxetine
for depression, 787
for insomnia, 499
for narcolepsy, 398
potentially inappropriate prescription of, 767

Flurazepam
for insomnia, 494, 496
potentially inappropriate prescription of, 764

Flutter, atrial, electrocardiography in, 908–909

Fluvoxamine, for insomnia, 499

Food pyramid, for persons over 70 years of age, 661

Foot, biomechanics of
in jumping, 247–249
in running, 245–247

Foot injuries
anatomic considerations in, 105–106
apophysitis, 148–150
examination in, **105–132**
arches, 108
body alignment, 107–108
forefoot, 127–129
gait, 109–111
hindfoot, 122–126
innervation, 115–116
midfoot, 126–127
shape, 109
shoe-wear patterns, 111–113
skin, 112, 114–115
toes, 127–129
fractures, 127–128, 143
Freiberg's infraction, 151
Iselin's disease, 150
jogger's foot, 127
Köhler's disease, 126, 150
Lisfranc's sprains, 126–127, 143–145
osteochondroses, 150–151
peroneal tendon, 153–155
plantar fasciitis, 155–156
Sever's disease, 149–150
stress fractures, 146–148
talar osteochondral lesions, 153
tarsal coalition, 151–152
turf toe, 156–157

Foot slap gait, in athletes, 109–110

Football, spear tackling in, cervical injury in, 233

Forefoot injuries, 127–129

Fracture(s). *See also specific bone.*
finger, 100–102
foot, 127–128, 143
groin region, 189–191
hand, 100–102
hip, in pediatric patients, 173–174
sacral, 220
stress. *See* Stress fractures.
toe, 129
vertebral, in athletes, 204–208, 212–213, 215, 218–219
wrist. *See* Wrist injuries, fractures.

Framingham Risk Score
in coronary artery disease, 932, 940, 1001–1004
in erectile dysfunction, 983, 986
in hyperlipidemia, 1034–1037

Franklin Cardiovascular Health Program, 867, 870

Freiberg's infraction, 151

Froment's sign, in hand injuries, 82

Functional Activities Questionnaire, 631–632, 835

Functional status, in nutritional assessment, 661–663

Functional urinary incontinence, 713

Furosemide, for congestive heart failure, 1124–1125

G

Gabapentin, for restless legs syndrome, 444, 576

Gait
assessment of
for fall risk, 688–690
in athletes, 109–111
screening instruments for, 841
in running, 245–247

Gamekeeper's (skier's) thumb, 87–88, 96–97

Gamma hydroxybutyrate, for narcolepsy, 398, 400

Gamma-aminobutyric acid receptor agonists, for insomnia, 383–384

Ganglia, wrist, 63

Gastroesophageal reflux disease, sleep disorders in, 524–527, 555

Gemfibrozil, for hyperlipidemia, 1049

Gender differences
in central sleep apnea, 365
in coronary artery disease risk, 964

Gene therapy, for hyperlipidemia, 1051

Generalized anxiety disorder, sleep disorders in, 545–546

Genetic factors
in delayed sleep phase circadian sleep disorders, 454
in depression, 779
in hypertension, 727
in narcolepsy, 394–397
in restless legs syndrome, 439–440

Genitofemoral nerve, entrapment of, 176–177

Geriatric assessment, incremental. *See* Incremental geriatric assessment.

Geriatric Depression Scale, 632, 844

Geriatric syndrome, 602

German Cardiovascular Prevention Study, 868

Get Up and Go Test, 627, 841

Glucose intolerance, in obstructive sleep apnea, 335

Glutethimide, for insomnia, 493, 497

Glycoprotein IIB/IIIa inhibitors, for coronary artery disease, 1068

Golfer's elbow, 7–8

Gout, coronary artery disease in, 972

Grafts, osteochondral, for knee osteochondral injuries, 267–268

Greenstick fractures, of distal radius, 48

Grief
depression in, 782
in end-of-life care, 824–825

Groin disruption, 179–180

Groin injuries, **163–183, 185–200**
acetabular labral tears, 191–193
adductor strains, 166–168
anatomic considerations in, 164–166
articular cartilage injuries of hip, 193–194
avascular necrosis of hip, 194–196
causes of, 163
differential diagnosis of, 163
epidemiology of, 163
femoral neck fractures, 189–191
femoral nerve entrapment, 175–176
genitofemoral nerve entrapment, 176–177
groin disruption, 179–180
high hamstring strains, 171
hip avulsion fractures and apophysitis, in pediatric patients, 173–174
hip disorders, 191–197
iliohypogastric nerve entrapment, 176–177
ilioinguinal nerve entrapment, 176–177
iliopsoas strains and bursitis, 169–171
ischial tuberosity avulsion, 171
lateral femoral nerve entrapment, 178
ligamentum teres, 194
nerve entrapment, 174–178
obturator nerve entrapment, 174–175

osteitis pubis, 186–188
posterior abdominal wall, 178–180
pubic ramus fractures, 189–191
pubic symphysis osteomyelitis, 188–189
rectus femoris strains, 172
sartorius strains, 172
snapping hip syndromes, 196–197
sports hernia, 178–179
strains, 166–173
stress fractures, 189–191

Growth hormone, as ergogenic aid, 286

Guanadrel, potentially inappropriate prescription of, 767

Guanethidine, potentially inappropriate prescription of, 767

Guyon's canal
anatomy of, 21–22, 24
ulnar nerve compression at, 65

H

Haddad syndrome, respiratory failure in, 552–553

Haglund's deformity, 121

Hallucinations, hypnagogic, in narcolepsy, 391

Hallux rigidus, 128

Hallux valgus, 128

Hamate
anatomy of, 21
fracture of, 56–57

Hammertoe, 128–129

Hamstring muscles
biomechanics of, 249–251
strains of, 171

Hand injuries, **71–89, 91–103**. *See also* Finger injuries.
anatomic considerations in, 72–73, 92–93
boxer's knuckle, 100
history of, 72–75, 91–92
misdiagnosis of, 71
physical examination in, 75–88, 93–95
Elson test in, 85–86
Finkelstein test in, 87
flexor digitorum profundus and superficialis testing in, 84–85
inspection in, 75, 77
ligamentous instability test in, 87–88
mallet finger test in, 85
neurovascular, 80–81, 94
palpation in, 77–78
range of motion assessment in, 78–80
strength testing in, 81–84
volar plate assessment in, 86
types of, 74

Head banging, in sleep, 421, 554

Health care triad, for dementia care, 672

Hearing, screening instruments for, 631, 831

Heart
disease of. *See* Cardiovascular disease; Coronary artery disease.
transplantation of, hyperlipidemia in, 1052

Heart block, electrocardiography in, 911–915

Heart failure, congestive. *See* Congestive heart failure.

Heart Protection Study, 1021, 1051

Heart rate, resting and peak, in cardiovascular disease prediction, 971

Heart sounds, in congestive heart failure, 1118

Heel rise test, in tibialis posterior dysfunction, 123–125

Hemarthrosis, in osteochondral injuries, 258–259

Hemoglobin, artificial increase of, for sports performance, 282–283

Hemorrhage, intracranial, in thrombolytic therapy, 1071

Heparin, for coronary artery disease, 1068, 1071

Herbal preparations
drug prescription precautions with, 769–770
for hyperlipidemia, 1044–1045
for insomnia, 498

Herbert classification, of scaphoid fractures, 50–51

Hernia, sports, 178–179

Hindfoot injuries, examination in, 122–126

Hip
acetabular labral tears of, 191–193
apophysitis of, in pediatric patients, 173–174
articular cartilage injuries of, 193–194
avascular necrosis of, 194–196

Hip (*continued*)
avulsion fractures of, in pediatric patients, 173–174
biomechanics of
in jumping, 247–249
in kicking, 249–251
in running, 246–247
snapping, 196–197

Hirschsprung's disease, respiratory failure with, 552–553

Hirudin, for coronary artery disease, 1068

Histamine antagonists, for narcolepsy, 400

Home health care, in dementia, 677

Home safety, assessment of, 636, 854. *See also* Incremental geriatric assessment.

Homeostasis, in sleep, 321

Homocysteine, elevated, coronary artery disease risk in, 968

Hormone replacement therapy, hypertension due to, 727

Hospice services, 814–815
for dementia, 678
in nursing facilities, 797

Human immunodeficiency virus infection, hyperlipidemia in, 1052

Human leukocyte antigens, in narcolepsy, 394–395

Huntington's disease, sleep disorders in, 540

Hydantoins, sleepiness due to, 503

Hydralazine
for congestive heart failure, 1125
for hypertension, 738

Hydrochlorothiazide
for congestive heart failure, 1125
for hypertension, 737–739, 1019–1020

Hydromorphone, for pain, in end-of-life care, 817

β-Hydroxy-β-methylbutyrate, as ergogenic aid, 287

Hydroxymethyl glutaric acid-co-reductase inhibitors (statins), for hyperlipidemia, 1045–1047

Hydroxyzine, potentially inappropriate prescription of, 766

Hyoscyamine, potentially inappropriate prescription of, 766

Hyperapolipoproteinemia, 1041

Hypercalcemia, electrocardiography in, 925

Hypercholesterolemia
familial, 1004–1005, 1037–1038
polygenic, 1005, 1034, 1037

Hypercoagulability, in obstructive sleep apnea, 512

Hyperhomocysteinemia, coronary artery disease risk in, 968

Hyperkalemia, electrocardiography in, 925

Hyperkeratosis, in foot, 114

Hyperlipidemia, **1027–1055**
antihypertensive drug-induced, 743–744
apolipoprotein B excess, 1041
atherogenic dyslipidemia, 1041–1042
atherosclerosis in, 1027–1028, 1030–1031, 1038–1039
broad band dyslipoproteinemia, 1041
coronary artery disease risk in, 965, 1034–1037
defining, 1034–1042
drug-resistant, 1050
familial combined, 1005, 1041
familial defective apolipoprotein B_{100}, 1005, 1038
familial high-density lipoprotein deficiency, 1042
familial hypercholesterolemia, 1037–1038
familial hypertriglyceridemia, 1040
familial hypoalphalipoproteinemia, 1042
hyperapolipoproteinemia, 1041
in diabetes mellitus, 1051–1052
in elderly persons, 1052–1053
in erectile dysfunction, 979
in heart transplantation, 1052
in HIV infection, 1052
in kidney transplantation, 1052
in pediatric patients, 1034
isolated low high-density lipoprotein cholesterol, 1042
lecithin cholesterol acyl transferase deficiency, 1042
lipid measurement in, 1033–1034
lipoprotein metabolism and pathophysiology in, 1028–1033
lipoprotein(a) excess, 1038–1039
low-density lipoprotein-dominant, 1034, 1037
mixed lipid disorder, 1040–1041
polygenic, 1005, 1034, 1037
secondary, 1043
severe hypertriglyceridemia, 1040
sitosterolemia, 1038
Tangiers disease, 1042

treatment of
follow-up, 1053
goals of, 1043
in coronary artery disease, 1064, 1073–1074
initiation of, 1043–1053
types of, 1027–1028

Hyperprolactinemia, erectile dysfunction in, 988

Hypersomnia. *See also* Narcolepsy; Sleepiness, excessive daytime, idiopathic, 402–404

Hypertension, **723–753, 1011–1025**
accelerated, 744
age-related factors in, 727–728
ambulatory blood pressure monitoring in, 1014–1016
blood pressure circadian variation and, 731
cardiovascular effects of, 728–731, 965
coronary artery disease in, 965
definition of, 1012–1013
epidemiology of, 1011
evaluation of, 731–735
in very elderly persons, 744–745
isolated systolic, 723–724, 728–729
labile, 734–735
obstructive sleep apnea and, 307, 334, 512–514
prevalence of, 723–726
risk factors for, 726–727, 1011–1012
secondary, 1014
treatment of, 735–745
1073, 1073–1074
after myocardial infarction, 1022
after stroke, 1022–1023
approach to, 1014
goals of, 1012–1014
in angina, 1021–1022
in diabetes mellitus, 741–742, 744, 1021
in heart failure, 1022
in kidney disease, 1021
in pediatric patients, 1023
in renal dysfunction, 741
lifestyle modifications in, 1016–1018
nonpharmacologic, 736
nonstandard, 1021–1023
pharmacologic, 736–741, 1018–1021
side effects of, 742–744
white-coat, 1015

Hypertension Optimal Treatment (HOT) Trial, 1013

Hypertriglyceridemia
familial, 1005, 1040
in pancreatitis, 1031
severe, 1040

Hyperuricemia, coronary artery disease in, 972

Hypnagogic hallucinations, in narcolepsy, 391

Hypnic jerks (sleep starts), 425–426

Hypnopompic phenomena, in narcolepsy, 390–391

Hypnotics, for insomnia, side effects of, 492–498

Hypoalphalipoproteinemia, 1041
familial, 1042

Hypocalcemia, electrocardiography in, 925

Hypocapnia, in central sleep apnea, 363–364

Hypocretin, deficiency of, in narcolepsy, 395–397, 558–559

Hypogonadism, erectile dysfunction in, 988

Hypokalemia, electrocardiography in, 923, 925

Hypomagnesemia, electrocardiography in, 925

Hypopnea, in sleep. *See also* Obstructive sleep apnea-hypopnea syndrome, definition of, 330

Hypothalamus
disorders of, narcolepsy in, 391–393
in sleep control, 535–536

Hypothenar eminence, palpation of, 78

Hypothyroidism, central sleep apnea in, 367

I

Ibutilide, for atrial fibrillation, 1094

Iliohypogastric nerve, entrapment of, 176–177

Ilioinguinal nerve, entrapment of, 176–177

Iliopsoas muscle
bursitis of, 169–171
strains of, 169–171
tendonitis of, 169–171

Imipramine, for insomnia, 498–499

Immune system, sleep deprivation effects on, 480

Immunizations, for elderly persons, 606

Implantable cardiac defibrillators, for coronary artery disease, 1075–1076

Incontinence, urinary. *See* Urinary incontinence.

Incremental geriatric assessment, **619–643**
care needs determined by, 638–639
first level (early signs of dysfunction), 621–630
monitoring care in, 640
second level (significance of functional impairments), 631–636
summary of, 641
third level (comprehensive), 636–638

Indiplon, for insomnia, 383, 494

Indomethacin, potentially inappropriate prescription of, 764

Infections, coronary artery disease risk in, 972

Information
on exercise, for fall prevention, 695
on urinary incontinence, 715

Informed consent
in nursing facilities, 804
in shared decision making, 646–647

Injuries, sports. *See* Sports injuries.

Injury, in falls, 683–685

Insomnia, **375–388**
behavioral, 553
causes of, 379–380
conditions associated with, 376–377
consequences of, 376–377
definition of, 376
diagnosis of, 377–379
drugs inducing, 492
epidemiology of, 375–376
in depression, 543
in elderly persons, 566–569
in pediatric patients, 553
mood disorders and, 385–386
overview of, 312–314
primary, 380
treatment of, 380–385, 492–498

Instrumental activities of daily living, assessment of, 661–663, 832–833. *See also* Incremental geriatric assessment.

Instruments, for screening, **829–854**
alcoholism, 849–850
caregiver burden, 851–853
clock drawing task, 834
cognitive function, 834–840
delirium criteria, 840
dementia criteria, 838–839
depression, 844–848
functional activities questionnaire, 835
gait, 841
hearing, 831
home safety, 854
instrumental activities of daily living, 832–833
mini-mental state examination, 836–837
mobility, 841–843
nutrition, 830

Insufficient sleep syndrome, 476–477

Insulin resistance, in obstructive sleep apnea, 335, 512

Internal fixation
of distal radial fracture, 48
of scaphoid fracture, 53–54

International Classification of Sleep Disorders, 377–378

International Olympic Committee, ergogenic aid policies of, 278

International Restless Legs Study Group criteria, 435–437

Interpersonal therapy, for depression, 786

Interphalangeal joints
anatomy of, 72
proximal, dislocation of, 101–102

Intersection syndrome, 38–40

Intervertebral disc disease
cervical, 238–239
lumbar, 215–218

Intracranial hemorrhage, in thrombolytic therapy, 1071

Intraventricular conduction delay, nonspecific, electrocardiography in, 921

Inversion stress test (talar tilt test), in ankle sprain, 118–119, 136–137

Ipratropium, for COPD, 524

Iron deficiency, restless legs syndrome in, 439, 441–442, 576

Iron overload, coronary artery disease risk in, 973

Irregular sleep–wake circadian rhythm sleep disorders, 459

Ischemic heart disease, heart failure in, 1116

Ischial tuberosity, avulsion of, 171

Iselin's disease, 150

Isoxsurpine, potentially inappropriate prescription of, 768

Isradipine, for hypertension, 739

J

Jactatio capitis nocturna, 554

Jersey finger, 84–85, 98–99

Jet lag, 460–462

Jogger's foot, 127

Jugular venous distention, in congestive heart failure, 1118

Jumping, biomechanics of, 247–249

Junctional rhythm, electrocardiography in, 910–911

Juvenile plantar dermatosis, 114

K

Kava, for insomnia, 498

Kegel exercises, for urinary incontinence, 709, 715–716

Keratolysis, pitted, in foot, 114

Ketorolac, potentially inappropriate prescription of, 767

Kicking, biomechanics of, 249–251

Kidney
- dysfunction/failure of
  - coronary artery disease in, 967
  - hypertension treatment in, 1021
  - sleep disorders in, 367, 527–529
- transplantation of, hyperlipidemia in, 1052

Kienböck's disease, 55–56

Klein-Levin syndrome, 559

Knee
- biomechanics of
  - in jumping, 249
  - in kicking, 249–251
- osteochondral injuries of
  - nonoperative management of, 264–265
  - physical examination in, 259–260
  - surgical treatment of, 265–270

Köhler's disease, 126, 150

L

Labile hypertension, 734–735

Lai Tai (sudden unexplained nocturnal death syndrome), 428–429

Landau-Kleffner syndrome, sleep disorders in, 555

Lateral epicondylitis (tennis elbow), 4–6

Lateral foot walking test, in ankle injuries, 110

Lateral pivot shift test, in posterolateral rotatory instability, 11

Lavage, for knee osteochondral injuries, 266

Laxatives, potentially inappropriate prescription of, 767

Lecithin cholesterol acyl transferase, 1030
- deficiency of, 1042

Left atrial enlargement, electrocardiography in, 903–904

Left bundle branch blocks, electrocardiography in, 918–919

Left ventricular hypertrophy
- coronary artery disease risk in, 966
- electrocardiography in, 915–916

Leg(s)
- nocturnal cramps of, 417, 440
- periodic movement of. *See* Periodic leg movement disorder.
- restless. *See* Restless legs syndrome.

Legal issues, in nursing facilities, 803–804

Leptin regulation, in obstructive sleep apnea, 335

Levodopa-carbidopa, for restless legs syndrome, 441–443, 576

Lewy body disease
- dementia with, sleep disorders in, 538
- rapid eye movement sleep behavior disorder in, 419

Life support, for cervical spinal injuries, 231–233

Lifestyle modifications
- for congestive heart failure, 1122
- for coronary artery disease prevention, **947–961,** 1073–1074
  - diet in, 953–955
  - exercise in, 950–953
  - multiple behavior interventions in, 955–957
  - smoking cessation in, 947–950
- for fall prevention, 695
- for hyperlipidemia, 1043–1044
- for hypertension, 1016–1018
- for obstructive sleep apnea, 345
- for restless legs syndrome, 441
- for urinary incontinence, 709–714

Ligamentous laxity, cervical, 237–238

Ligamentum teres, injury of, 194

Light
in circadian rhythm maintenance, 449–450, 452
therapy with
for advanced sleep phase disorders, 457, 578
for circadian rhythm sleep disorders, 506
for delayed sleep phase disorders, 455
for irregular sleep–wake disorders, 459
for jet lag, 460–462
for nonentrained sleep phase disorders, 459
for shift work sleep disorders, 464

Lighting, for sleep deprivation, 486

Lipid disorders. *See* Hyperlipidemia.

Lipoprotein(a), excess of, 969–970, 1038–1039

Lipoprotein(s)
disorders of. *See* Hyperlipidemia.
high-density, 1030
deficiency of, 1042
in hypoalphalipoproteinemia, 1042
isolated low levels of, 1042
intermediate-density, 1029
low-density, 1029–1030
apheresis of, 105–1051
excess of, 1034, 1037
in atherosclerosis, 1030–1031
oxidation of, 1031
receptor for, defects of, 1037–1038
measurement of, 1033–1034
metabolism of, 1028–1030
receptors for, defects of, 1037–1038, 1050
types of, 1028–1030
very low-density, 1029, 1031, 1041

Lipoprotein lipase, 1028
deficiency of, 1005

Lisfranc joint, sprain of, 126–127, 143–145

Lisinopril, for hypertension, 739, 1020, 1022–1023

Lithium, for depression, 787

Lo standards, for determination of capacity, 650

Lorazepam, potentially inappropriate prescription of, 765

Losartan, for hypertension, 739

Lunate
anatomy of, 19
avascular necrosis of, 55–56
fracture of, 55–56

Lunate dissociation, complete, 59

Lunotriquetral ligaments, injury of, 60–62

Lysine, as ergogenic aid, 287

M

Maculopathy, age-related, in hypertension, 729

Magnetic resonance arthrography
in acetabular labral tears, 192
in osteochondral injuries, 262–263

Magnetic resonance imaging
functional, in restless legs syndrome, 438–439
in adductor strains, 167
in depression, 782
in femoral neck stress fracture, 190
in foot fractures, 146
in hip articular cartilage injuries, 193
in osteochondral injuries, 261–262
in rapid eye movement sleep behavior disorder, 420
in scaphoid fracture, 51
in scapholunate dissociation, 59–60
in spondylolysis, 206
in symphysis pubis, 187
in tarsal coalition, 152
in triangular fibrocartilage complex injury, 45
in ulnar collateral ligament injury, 9
in wrist injuries, 32

Maintenance of Wakefulness Test, 310

Make Early Diagnosis to Prevent Early Death (MEPED), in coronary artery disease, 1005

MALES (Men's Attitudes to Life Events and Sexuality) Study, 978

Mallet finger, 85, 97–98

Mallet toe, 128

Malnutrition
adverse effects of, 664–665
interventions for, 666–667

Mandible, appliances for, for obstructive sleep apnea, 349, 573

Manipulation
for back pain, 222
for cervical sprain, 238

Maprotiline, for insomnia, 499

March fractures. *See* Stress fractures.

Marrow-stimulating techniques, for knee osteochondral injuries, 266–267

Massachusetts Male Aging Study (MMAS), 978, 982

Masticatory muscle activity, rhythmic, in bruxism, 422–423

Maze procedure, for atrial fibrillation, 1102–1103

Mechanical ventilation
for central sleep apnea, 369–371
for obstructive sleep apnea, 513–514

Medial epicondylitis (golfer's elbow), 7–8

Medial foot walking test, in ankle injuries, 110

Median nerve
at elbow
anatomy of, 2
compression of, in pronator syndrome, 3–4
at wrist
anatomy of, 24
compression of, 64–65
examination of, 28–29

Medical Research Council study, on hypertension, 740

Medications. *See* Drug(s); *specific drugs*.

Melatonin
for advanced sleep phase disorders, 456, 578–579
for delayed sleep phase disorders, 455
for insomnia, 495, 498
for irregular sleep-wake disorders, 459
for jet lag, 462
for nonentrained sleep phase disorders, 459
for shift work sleep disorders, 464
in circadian rhythm control, 451–452
safety issues with, 465–466
side effects of, 506
sleep deprivation effects on, 480–481

Memory deficits, in sleep deprivation, 478–479

Menopause, sleep disorders after, 564

Men's Attitudes to Life Events and Sexuality (MALES) Study, 978

Mental status testing, determination of capacity and, 650

MEPED (Make Early Diagnosis to Prevent Early Death), in coronary artery disease, 1005

Meperidine, potentially inappropriate prescription of, 767

Meprobamate, potentially inappropriate prescription of, 765

Mesoridazine, potentially inappropriate prescription of, 768

Metabolic syndrome, coronary artery disease and, 967, 1074

Metacarpal bones
anatomy of, 72
fifth, fracture of, 100–101
palpation of, 77

Metacarpophalangeal joints
anatomy of, 72
injury of, boxer's knuckle as, 100
inspection of, 75
palpation of, 77
range of motion assessment of, 79

Metatarsal bones
fifth, apophysitis of, 150
fractures of, 127–128
head of, infarction of, 151

Metatarsophalangeal joint
first, sprain of, 156–157
injuries of, 128

Methandrostenolone, as ergogenic aid, 280

Methaqualone, for insomnia, 493, 497

Methyldopa
for hypertension, 738, 742
potentially inappropriate prescription of, 766

Methylphenidate
for depression, 787–788
for narcolepsy, 397–399, 500

Methyltestosterone, potentially inappropriate prescription of, 768

Methyprylon, for insomnia, 493, 497

Metoclopramide, for nausea and vomiting, in end-of-life care, 818

Metolazone, for congestive heart failure, 1125

Metoprolol
for atrial fibrillation, 1091
for congestive heart failure, 1123–1124
for coronary artery disease, 1064, 1068
for hypertension, 738–740

MHHP (Minnesota Heart Health Program), 868, 870–871

Michigan alcoholism screening test, 849

Midfoot injuries, examination in, 126–127

Mineral oil, potentially inappropriate prescription of, 768

Mini Nutritional Assessment, 665, 669

Mini-Cog test, 627

Mini-Mental State Examination, 631–632, 836–837

Minnesota Heart Health Program (MHHP), 868, 871

Mirtazapine
for depression, 787–788, 821
for insomnia, 498–499

Mitral valve
disorders of, electrocardiography in, 903–904
prolapse of, 921, 1110, 1113
regurgitation of, 1110, 1112–1113
stenosis of, 1110, 1113–1114

Mixed lipid disorder, 1040–1041

Mixed urinary incontinence, 708, 712

MMAS (Massachusetts Male Aging Study), 978, 982

Mobility, assessment of, 841–843

Mobitz type heart blocks, electrocardiography in, 912–914

Modanifil
for narcolepsy, 398–399, 500, 559
for sleep deprivation, 486–487

Monoamine oxidase inhibitors
for insomnia, 499
sleepiness due to, 503

Mood disorders
insomnia and, 385–386
sleep disorders in, 542–544

Morphine
for coronary artery disease, 1069
for pain, in end-of-life care, 817

Morton's neuroma, 128

Motor vehicle accidents, due to sleep deprivation, 481

Multidetector row computed tomography, in coronary artery disease, 938–939, 941

Multifascicular block, electrocardiography in, 920

Multifocal atrial tachycardia, electrocardiography in, 909–910

Multiple Sleep Latency Test, 309–310
in elderly persons, 563–564
in narcolepsy, 393–394, 401
in sleep deprivation, 484

Multiple system atrophy, sleep disorders in, 418–419, 538–539

Multiple wavelet model, of atrial fibrillation, 1085–1086

Murmurs, heart, in valvular disease, 1109–1113

Muscle cramps, nocturnal, 417, 440

Muscle relaxants
potentially inappropriate prescription of, 764
sleepiness due to, 502

Muscle strains, in back, 213–214, 216

Myasthenia gravis, sleep disorders in, 540

Myocardial infarction
congestive heart failure in, 1118
electrocardiography in, 922–924, 926
hypertension after, treatment of, 1022
non-ST elevation, treatment of, 1067–1070
risk assessment in, 1066–1067
ST elevation, 1070–1071
treatment of, 1067–1070, 1075

Myocardial ischemia
erectile dysfunction in, 983–984
heart failure in, 1116

Myopathy, statin-related, 1047

Myotonic dystrophy, sleep disorders in, 540

N

Nail disorders
in foot, 114–115
in hand, 102

Napping
in elderly persons, 563–564
in narcolepsy, 390, 401
in sleep deprivation, 485–486

Naproxen, potentially inappropriate prescription of, 767

Narcolepsy, **389–413**
cataplexy and, 390, 394, 399–402
clinical features of, 389–390

definition of, 558
diagnosis of, 393–394
differential diagnosis of, 402–405
genetic factors in, 394–397
idiopathic, 389
in pediatric patients, 558–559
overview of, 312
pathophysiology of, 390–393
prevalence of, 389
rapid eye movement sleep behavior disorder with, 419
treatment of, 397–399
versus idiopathic hypersomnia, 402–404

Nasal congestion, obstructive sleep apnea in, 332

National Cholesterol Education Program, coronary artery disease risk score of, 1000, 1003

Natural killer cells, sleep deprivation effects on, 480

Nausea and vomiting, in end-of-life care, 818–819

Navicular, osteochondrosis of (Köhler's disease), 126, 150

Neck injuries. *See* Cervical injuries.

Nefazodone
for depression, 787
for insomnia, 498–499

Nerve conduction studies
in carpal tunnel syndrome, 65
in wrist injuries, 32

Nervous system, function of, in sleep, 324

Neurocristopathy syndrome, 552–553

Neurologic disorders
erectile dysfunction in, 988
sleep disorders with, 536–542

Neurolysis, in obturator nerve entrapment, 175

Neuroma, Morton's, 128

Neurontin, for restless legs syndrome, 442

Neuropraxia
at wrist, 63–64
cervical spinal cord, 234–236
in stingers, 234

Neurotransmitters, abnormalities of, narcolepsy in, 391–393

New York Heart Association, functional classification of, 1121

Niacin, for hyperlipidemia, 1049–1050

Nicotine replacement therapy, for smoking cessation, 949, 1073

Nicotinic acid, for hyperlipidemia, 1049–1050

Nifedipine
for coronary artery disease, 1069
potentially inappropriate prescription of, 768

Nightmares, 416–417
in pediatric patients, 554
overview of, 314–315

Nitrates
for congestive heart failure, 1125
for coronary artery disease, 1065, 1069

Nitrendipine, for hypertension, 738

Nitric oxide, in erectile function, 980–982

Nitrofurantoin, potentially inappropriate prescription of, 768

Nitroglycerin, for coronary artery disease, 1069, 1073

Nocturnal leg cramps, 417, 440

Nocturnal paroxysmal dystonia, 417–418

Nocturnal urinary incontinence, 700, 714

Nonentrained circadian rhythm sleep disorders, 458–459

Noninvasive positive pressure ventilation, for central sleep apnea, 369–371

Nonlaita (sudden unexplained nocturnal death syndrome), 428–429

Non-rapid eye movement sleep. *See* Sleep, non–rapid eye movement (NREM).

Nonsteroidal anti-inflammatory drugs
for back pain, 223
for lateral epicondylitis, 5
for medial epicondylitis, 8
for pronator syndrome, 4
for radial tunnel syndrome, 6
for triceps tendon injury, 13
hypertension due to, 726
potentially inappropriate prescription of, 767

Nordic Diltiazem Study, 739

Norepinephrine, elevated
in congestive heart failure, 1116–1117
in sleep deprivation, 479

NORSJO Project, 868, 872

North Karelia Project, 866–867, 870, 877–878

Nortriptyline
for depression, 787–788
for insomnia, 498–499

NREM sleep. *See* Sleep, non–rapid eye movement (NREM).

Nuclear perfusion imaging, in coronary artery disease, 935–937

Nursing facilities, **793–810**
demographics of, 795–796
historical background of, 794–795
hospice care in, 797, 815
long-term care in, 797
overview of
physician roles in
administrative, 804–806
attending, 799–803
ethical issues in, 803–804
in interdisciplinary team, 806–809
visits, 799–802
regulations on, 797–799
rehabilitation in, 797
skilled care in, 797
staff of, 795
types of care in, 796–799

Nutrition, **659–669**. *See also* Diet.
assessment of, 621–624, 626, 631, 830
functional status in, 661–663
in disease states, 663–664
tools for, 661, 665–666, 668–669
in end-of-life care, 819
in normal aging, 659–660
problems with
adverse effects of, 664–665
interventions for, 666–667

O

Obesity
cardiovascular disease in, 966
hypertension in, 727, 733–734, 1017
obstructive sleep apnea in, 332, 341

Obesity hypoventilation syndrome, 331

Obsessive-compulsive disorder, sleep disorders in, 545

Obstructive sleep apnea-hypopnea syndrome, **329–359**
cardiovascular disease in, 333–335, 511–515
Cheyne-Stokes respiration in, 515–518
clinical features of, 556
consequences of, 333–336
COPD with, 522–523
diagnosis of, 337–344
epidemiology of, 336–337
hypertension and, 307, 334, 512–514
in Alzheimer's disease, 537
in elderly persons, 569–574
in pediatric patients, 555–557
insomnia in, 492
kidney failure with, 528–529
laboratory studies in, 341
metabolic consequences of, 335
obstruction sites in, 350
overview of, 306–309
pathophysiology of, 331–333, 511–512
periodic leg movement disorder with, 440–441
perioperative risk in, 336
polysomnography in, 343–344
prediction formulae for, 341–342
pulse oximetry in, 342
risk factors for, 344–353, 556
terminology of, 329–331
treatment of, 344–351, 499–500, 513–515, 572–574
versus narcolepsy, 404–405

Obturator nerve, entrapment of, 174–175

Occupational therapy, progress in, review of, 634

Office systems, for heart disease prevention, **883–900**
goals of, 895–896
implementation of, 893–898
management in, 886–893
monitoring in, 893
protocols for, 895
reviewing and revising services in, 894, 897–898
screening in, 884–885

Older adults. *See* Elderly persons.

Omnibus Budget Reconciliation Act of 1987, nursing facility provisions of, 794–795, 797–799

Ondine curse (central hypoventilation syndrome), 552–553

Onychocryptosis, 114

Onychodystrophy, in athletes, 114

Onychogryphosis, in athletes, 114–115

Onychomycosis, in athletes, 115

Opioids
agonists of, sleepiness due to, 502–503
constipation due to, 820
for dyspnea, in end-of-life care, 818
for pain, in end-of-life care, 817
for restless legs syndrome, 444

Oral appliances, for obstructive sleep apnea, 349

Orexins, deficiency of, in narcolepsy, 395–397

Ornithine, as ergogenic aid, 287

Orphenadrine, potentially inappropriate prescription of, 767

Orthodromic atrioventricular re-entry tachycardia, electrocardiography in, 906–907

Osler's maneuver, in blood pressure measurement, 732

Osteitis pubis, 186–188

Osteochondral injuries, **253–276**
arthroscopy in, 263–264, 266, 272
capitellum, 273
cartilage composition and function and, 253–255
clinical presentation of, 258–264
definitions of, 253
foot, 150–151
history of, 258–259
imaging in, 260–264
knee, 259–260, 264–271
navicular (Köhler's disease), 126, 150
pathophysiology of, 255–258
physical examination in, 259–260
talus, 121–122, 153, 260, 270, 272
treatment of, 264–273
capitellum, 273
knee, 265–271
nonoperative, 264–265
options for, 264
surgical, 265–273
talus, 270, 272

Osteochondritis dissecans
ankle, 121–122, 153
definition of, 253
talus, 270, 272

Osteomyelitis, of pubic symphysis, 188–189

Ottawa ankle rules, 118, 137, 140

Overflow urinary incontinence, 712

Over-the-counter medications, drug prescription precautions with, 769–770

Overuse injuries. *See also* Stress fractures.
of wrist, 36

Oxaprozin, potentially inappropriate prescription of, 767

Oxazepam, potentially inappropriate prescription of, 765

Oxidation, of low-density lipoproteins, 1031

Oxidative stress, in obstructive sleep apnea, 512

Oxycodone
for pain, in end-of-life care, 817
for restless legs syndrome, 442, 444

Oxygen therapy
for central sleep apnea, 370
for Cheyne-Stokes respiration, 517–518
for chronic obstructive pulmonary disease, 522
for dyspnea, in end-of-life care, 818

P

P wave characteristics, in electrocardiography, 901–910

PACE + trial, for coronary artery disease prevention, 956

Pacemakers, for atrial fibrillation, 1102

Pain
chest. *See* Angina; Chest pain.
management of, in end-of-life care, 816–818

Palate, surgery on, for obstructive sleep apnea, 350

Palliative care, end-of-life. *See* End-of-life care.

Pancreatitis, hypertriglyceridemia in, 1031

Panic disorder, sleep disorders in, 544

Paralysis, sleep, 424–425
in narcolepsy, 390–391
in pediatric patients, 554

Parasomnias, **415–434,** 501, 505
bruxism, 314, 421–424
classification of, 415
confusional arousals, 314–315, 416
in elderly persons, 579–583
in pediatric patients, 553–555
nightmares, 314–315, 416–417, 554
nocturnal leg cramps, 417, 440
nocturnal paroxysmal dystonia, 417–418
overlap syndrome, 419
overview of, 314–315
pathogenesis of, 415–416
rapid eye movement sleep behavior disorder, 418–420, 554
rapid eye movement sleep–related sinus arrest, 420–421
rhythmic movement disorder, 421
sleep paralysis, 390–391, 424–425, 554

Parasomnias (*continued*)
sleep starts, 425–426
sleep talking, 426
sleep terrors, 314–315, 426–427, 554
sleep-related abnormal swallowing syndrome, 425
sleep-related painful erections, 425
sleepwalking, 314–315, 427–428, 553–554
sudden unexplained nocturnal death syndrome, 428–429

Parkinson's disease
sleep disorders in, 418–419, 539–540
treatment of, sleepiness due to, 502

Paronychia, 114

Paroxetine
for depression, 787
for insomnia, 498–499

Paroxysmal nocturnal dyspnea, in congestive heart failure, 1118

Pars interarticularis, spondylolysis at, 204–208

PATCH (Planned Approach to Community Health), 874

Patient rights, in nursing facilities, 803–804

Pavor nocturnes, 554

Pawtucket Heart Health Program (PHHP), 868, 871

PBBP (Pawtucket Heart Health Program), 868, 870–871

Pediatric patients
back pain in, 202–214
hip avulsion fractures and apophysitis in, 173–174
hyperlipidemia in, 1034
hypertension in, 1023
normal sleep in, 549–551
sleep disorders in, **549–562**
behavioral insomnias of childhood, 553
central hypoventilation syndrome, 552–553
in medical disorders, 554
in psychiatric disorders, 554
Klein-Levin syndrome, 559
narcolepsy, 558–559
parasomnias, 553–554
periodic limb movement disorder, 557
public health issues in, 559–560
restless legs syndrome, 440, 557
rhythmic movement disorder, 421
sleep apnea, 554–557
sleep phase delay, 557–558
sudden infant death syndrome, 552

Pelvic floor exercises, for urinary incontinence, 709, 715–716

Pemoline, for narcolepsy, 397–399, 500

Pentazocine, potentially inappropriate prescription of, 764

Performance enhancers, for sports. *See* Ergogenic aids.

Performance-Oriented Mobility Assessment, 690, 842–843

Pergolide, for restless legs syndrome, 442, 576

Periodic leg movement disorder, 312
clinical features of, 435, 437
in elderly persons, 574–577
in kidney failure, 528–529
in pediatric patients, 557
obstructive sleep apnea with, 440–441
treatment of, side effects of, 505–506

Periodontal disease, coronary artery disease risk in, 972

Peroneal nerves
deep, entrapment of, 126
evaluation of, in injuries, 115

Peroneal tendons
injuries of, 153–155
tendonitis of, 124, 154

Peyronie's disease, erectile dysfunction in, 988

Phalanges, fractures of, 101

Phalen's sign
in carpal tunnel syndrome, 64
in tarsal tunnel syndrome, 125
in wrist injuries, 28

Pharynx, collapsibility of, in obstructive sleep apnea, 331

Phenelzine
for depression, 787
for insomnia, 499

Phenobarbital, for insomnia, 497

Phenothiazines, for dyspnea, in end-of-life care, 818

Phototherapy. *See* Light, therapy with.

Physeal fractures, of distal radius, 48

Physical activity. *See* Exercise.

Physical function, screening tests for, 627–628

Physical therapy
for adductor strains, 167–168
for lateral epicondylitis, 5
for medial epicondylitis, 8
for radial tunnel syndrome, 6
for triceps tendon injury, 13
progress in, review of, 634

Piano key sign, in distal radioulnar joint instability, 44

Pindolol, for hypertension, 740

Pisiform
anatomy of, 20
fracture of, 57–58

Pitted keratolysis, in foot, 114

Pivot shift test, lateral, in posterolateral rotatory instability, 11

Planned Approach to Community Health (PATCH), 874

Plant sterols, accumulation of, in sitosterolemia, 1038

Plantar fasciitis, 125–126, 155–156

Plantar nerves
entrapment of, 125, 127
evaluation of, in injuries, 115

Plantar warts, 112

P-mitrale (left atrial enlargement), electrocardiography in, 903–904

Pokkuri (sudden unexplained nocturnal death syndrome), 428–429

Polysomnography
in delayed sleep phase circadian sleep disorders, 455
in depression, 543
in kidney failure, 527
in narcolepsy, 393–394
in obstructive sleep apnea, 343–344, 570
in sleepiness, 311

Positive pressure therapy
for asthma, 520–521
for central sleep apnea, 369–370
for Cheyne-Stokes respiration, 517–518
for obstructive sleep apnea, 346–349, 513–514, 573

Positron emission tomography, in restless legs syndrome, 438

Posterior elbow impingement, 13–15

Posterior interosseous nerve syndrome, 7
at elbow, 5, 7
at wrist, 65

Posterolateral rotatory instability, of elbow, 11–12

Post-traumatic stress disorder, sleep disorders in, 545

Postvoid residual volume, in urinary incontinence, 708

Potassium, for hypertension, 736

potentially inappropriate prescription of, 1095

P-pulmonale (right atrial enlargement), electrocardiography in, 903

PR interval characteristics, in electrocardiography, 910–915

Pramipexole, for restless legs syndrome, 442–443, 576

Pre-excitation syndromes, electrocardiography in, 911

Prehypertension
definition of, 1012
lifestyle modifications for, 1016–1018
treatment of, 1014

Premature atrial complex, electrocardiography in, 905

Premature ventricular contraction, electrocardiography in, 921

Prescriptions, for elderly persons. *See* Drug(s), prescribing, for elderly persons.

Press test, in triangular fibrocartilage complex injury, 44

Pressure ulcers, in end-of-life care, 820

Prinzmetal's angina, 1064

Procainamide, for atrial fibrillation, 1094–1095

PROGRESS trial, for stroke prevention, 1022–1023

Progressive supranuclear palsy, sleep disorders in, 538

Promethazine, potentially inappropriate prescription of, 766

Pronation, evaluation of, 111

Pronator syndrome, of elbow, 3–4

Propafenone, for atrial fibrillation, 1094–1095, 1101

Propantheline, potentially inappropriate prescription of, 766

Propoxyphene
for restless legs syndrome, 442, 444, 576
potentially inappropriate prescription of, 764

Propranolol, for atrial fibrillation, 1091

Proprioception, evaluation of, in ankle sprain, 119

Prostate, benign hypertrophy of, erectile dysfunction in, 988

Protein requirements, for elderly persons, 660

Protriptyline
for insomnia, 499
for narcolepsy, 398

Pseudoboutonnière deformity, 86, 99

Pseudohypertension, 732

Psychiatric disorders. *See also specific disorders.*
sleep disorders with, 324, 542–547

Psychologic function, assessment of, 628–629

Psychosocial factors
in coronary artery disease, 967
in depression, 779
in end-of-life care, 820–821

Psychotherapy
for depression, 786
for insomnia, 382

Pubic ramus, stress fractures of, 189–191

Pubic symphysis
inflammation of, 186–188
osteomyelitis of, 188–189

Pulse oximetry, in obstructive sleep apnea, 342

Pump bump, 121–122

Q

Q wave characteristics, in electrocardiography, 922–923

QRS interval characteristics, in electrocardiography, 915–922

QT interval characteristics, in electrocardiography, 923

Quadriceps muscles, biomechanics of, 249–251

Quality of life
in insomnia, 377
with urinary incontinence, 700

Quinidine, for atrial fibrillation, 1095

Quinine, for nocturnal leg cramps, 417

Quitlines, for smoking cessation, 949

R

RACE (RAte Control versus Electrical cardioversion), for atrial fibrillation, 1097–1098

Radial artery
at elbow, anatomy of, 3
in hand, examination of, 81

Radial nerve
at elbow
anatomy of, 2–3
compression of, 5–6
at wrist
anatomy of, 24
compression of, 65–66
in hand, examination of, 80

Radial tunnel syndrome, 5–6

Radiography
in ankle sprains, 118, 137
in carpal instability, 59–62
in cubital tunnel syndrome, 10
in distal radioulnar joint instability, 43
in hip avascular necrosis, 195
in Kienböck's disease, 56
in Lisfranc joint injury, 145
in osteitis pubis, 187
in osteochondral injuries, 260
in posterior elbow impingement, 14
in posterolateral rotatory instability, 11
in scaphoid fracture, 50
in Scheuermann's disease, 210
in spondylolysis, 205–206
in tarsal coalition, 152
in triangular fibrocartilage complex injury, 45
in ulnar collateral ligament injury, 9
in vertebral fractures, 212–213
in wrist injuries, 30–31, 36–37, 46–48

Radionuclide bone scan. *See* Bone scan.

Radionuclide studies, in coronary artery disease, 935–937

Radioulnar joint, distal
anatomy of, 17–18
instability of, 42–44

Radius
fracture of, at wrist, 45–49
styloid process of, palpation of, 77

Rales, pulmonary, in congestive heart failure, 1118

Ramipril, for coronary artery disease, 1065

Range of motion assessment
of hand, 78–80, 94
of wrist, 26

Rapid eye movement sleep. *See* Sleep, rapid eye movement (REM).

Rapid eye movement sleep behavior disorder, 418–420
in elderly persons, 579–583
in pediatric patients, 554

Rapid eye movement sleep–related sinus arrest, 420–421

Rate control, for atrial fibrillation, 1096–1099, 1102

RAte Control versus Electrical cardioversion (RACE), for atrial fibrillation, 1097–1098

Reagan's test, in wrist injuries, 29

Rectus femoris muscle, strains of, 172–173

Red yeast rice, for hyperlipidemia, 1044–1045

Rehabilitation, in nursing facilities, 797

Relationship issues, erectile dysfunction in, 988

REM sleep. *See* Sleep, rapid eye movement (REM).

Reserpine
for hypertension, 738
potentially inappropriate prescription of, 766

Resistance training, for fall prevention, 691–692

Respiratory dysfunction, in end-of-life care, 818, 823

Respiratory effort-related arousals, 330

Respiratory system
sedative effects on, 505
sleep disorders and, 518–524

Restless legs syndrome, **435–448**
clinical features of, 435–437
diagnosis of, 440–441
epidemiology of, 437–438
genetic factors in, 439–440
in elderly persons, 574–577
in kidney failure, 528–529
in pediatric patients, 440, 557
overview of, 312
pathophysiology of, 438–439
treatment of, 441–444, 505–506, 576–577

Restraints, in nursing facilities, 803

Resuscitation, in cervical spinal injuries, 231–233

Reteplase, for coronary artery disease, 1071

Revascularization, coronary, 1065–1066, 1069–1070

Rheumatic heart disease, mitral stenosis in, 1113

Rheumatoid arthritis
athletic participation with, 240
cardiovascular disease in, 971

Rhythm control, for atrial fibrillation, 1096–1098, 1100–1103

Rhythmic masticatory muscle activity, in bruxism, 422–423

Rhythmic movement disorder, 421

Right atrial enlargement, electrocardiography in, 903

Right bundle branch blocks, electrocardiography in, 916–918

Right ventricular hypertrophy, electrocardiography in, 916

Rocking, body, in pediatric patients, 554

Romberg test, in ankle sprain, 119

Romhilt-Estes score, in left ventricular hypertrophy, 916

Ropinirole, for restless legs syndrome, 442–443, 576

Rotterdam Study, for hypertension, 739

Running, biomechanics of, 245–247

S

Sacroiliitis, in athletes, 220

Sacrum, stress fracture of, 220

Safety
home, assessment of, 636, 854
in dementia, 616

Salmeterol, for nocturnal asthma, 521

SALSA mnemonic, for depression screening, 629

Sartorius muscle, strain of, 172

Scaphoid
anatomy of, 18
fracture of, 49–55

Scaphoid shift maneuver, in wrist injuries, 29, 59

Scapholunate angle, in carpal instability, 61–62

Scapholunate dissociation, 59–60

Scapholunate joint, examination of, 26–27

Scapholunate ligaments, injury of, 59

SCCDPP (South Carolina Cardiovascular Disease Prevention), 869, 872–873

Scheuermann's disease, in athletes, 203, 210

Schizophrenia, sleep disorders in, 546

Schmorl's nodes, in Scheuermann's disease, 210

Scintigraphy, bone. *See* Bone scan.

Scoliosis, in athletes, 211–212

Scotty dog sign, in spondylolysis, 206

Screening instruments. *See* Instruments, for screening.

SEA-squared model, for drug prescribing, 772–773

"Second ankle bone," 126–127

Sedatives, for insomnia, side effects of, 492–498

Seizures, in nocturnal paroxysmal dystonia, 417–418

Selective serotonin reuptake inhibitors
for depression, 787–788
for insomnia, 499
for narcolepsy, 400
sleepiness due to, 503

Selegiline, for narcolepsy, 400

Sensations, in restless legs syndrome, 435–437

Sertraline
for depression, 787
for insomnia, 499

Sever's disease, 149–150

Sexual dysfunction. *See also* Erectile dysfunction.
in antihypertensive therapy, 743

Shared decision making, **645–658**
advance care planning and, 647–648
autonomy and, 645–646, 648–649
cultural issues in, 656–658
description of, 645–646
determination of capacity in, 649–650
example of, 652–654
for patients lacking capacity, 651–652
informed consent and, 646–647
respect for dignity in, 654–656

SHEP (Systolic Hypertension in the Elderly Program) trial, 730, 738, 742, 1013

Shift work
circadian rhythm sleep disorders in, 462–465
sleep deprivation and, 482, 484–486

Shoe wear patterns, in foot and ankle injuries, 111–113

Sick sinus syndrome, electrocardiography in, 905

SIDS (sudden infant death syndrome), 552

Single photon emission computed tomography
in coronary artery disease, 936–937
in restless legs syndrome, 438
in spondylolysis, 206

Single-leg heel raise test, in ankle injury, 110

Sinus arrest, rapid eye movement sleep–related, 420–421

Sinus arrhythmia, electrocardiography in, 905

Sinus tachycardia, electrocardiography in, 6, 906

Sitosterolemia, 1038

Skier's thumb, 87–88, 96–97

Skilled nursing care, in nursing facilities, 797

Skin care, in end-of-life care, 820

Skin disorders, of foot, 112, 114–115

Sleep
age-related changes in, 563–564
architecture of, 549–551
arousal from, 306
blood pressure changes in, 731
function of, 308–309
medical disorders and, **511–533**
cardiovascular, 333–335, 511–518
gastroesophageal reflux disease, 524–527, 555
in pediatric patients, 555

renal, 527–529
respiratory, 518–524
neurologic systems influencing, 535–536
non–rapid eye movement (NREM), 306, 309
central sleep apnea in, 366
in Alzheimer's disease, 537
in parasomnias, 553–554
in pediatric patients, 550–551
parasomnias related to, 415–417, 424, 428
ontogeny of, 550–551
rapid eye movement (REM), 306, 309
central sleep apnea in, 366
hypothalamic control of, 391–393
in Alzheimer's disease, 537
in dementia with Lewy bodies, 538
in depression, 543
in parasomnias, 501, 553–554
in pediatric patients, 549–551
in post-traumatic stress disorder, 545
in progressive supranuclear palsy, 538
in schizophrenia, 546
parasomnias related to, 415–428
sleep-onset, 394–395, 404–405
stimulant effects on, 500–501
urinary incontinence during, 700, 714

Sleep apnea. *See also* Central sleep apnea; Obstructive sleep apnea-hypopnea syndrome.
as stroke risk factor, 540–541
hypertension in, 727
screening for, 315

Sleep behavioral disorder, 314

Sleep deprivation, **475–490**
behavioral consequences of, 477–479
catastrophic accidents related to, 481–482
clinical assessment of, 483–484
consequences of, 477–482
definition of, 475–476
interventions for, 484–487
motor vehicle accidents related to, 481
overview of, 309
physiological consequences of, 479–481
prevalence of, 475–477
qualitative, 476
quantitative, 475–476
shift work and, 482, 484–486

Sleep diary, 483

Sleep disorders
central sleep apnea, **361–374**
circadian rhythm, 314–315, **449–473**
classification of, 305
evaluation of, sleep process matrix for, **319–328**
excessive somnolence, **389–413**
in elderly persons, **563–586**
in neurologic disorders, 536–542
in pediatric patients. *See* Pediatric patients, sleep disorders in.
in psychiatric disorders, 542–547
insomnia. *See* Insomnia.
narcolepsy. *See* Narcolepsy.
overview of, **305–318**
parasomnias. *See* Parasomnias.
periodic leg movement disorder. *See* Periodic leg movement disorder.
restless legs syndrome. *See* Restless Legs syndrome.
sleep apnea. *See* Central sleep apnea; Obstructive sleep apnea-hypopnea syndrome.
sleep deprivation, 309, **475–490**
treatment of. *See also specific disorders.*
medications for, **491–509**
types of, 491

Sleep drunkenness, 402–404

Sleep hygiene
for insomnia, 313, 381, 568–569
for shift workers, 485

Sleep hypoventilation syndrome, 331

Sleep latency, in sleep deprivation, 477–478

Sleep paralysis, 424–425
in narcolepsy, 390–391
in pediatric patients, 554

Sleep process matrix, for sleep disorder evaluation, **319–328**
alcohol effects in, 325–326
behavioral/psychiatric processes in, 324–325
cardiorespiratory processes in, 323–324
case study using, 325, 327–328
chart for, 325–327
circadian rhythms in, 320–321
developmental changes in, 322–323
drug effects in, 325–326
homeostatic processes in, 321
neurologic processes in, 324
ultradian rhythms in, 321

Sleep restriction therapy, for insomnia, 381–382, 569

Sleep starts, 425–426

Sleep talking, 426, 554

Sleep terrors, 314–315, 426–427

Sleep-disordered breathing. *See also* Central sleep apnea; Obstructive sleep apnea-hypopnea syndrome.
in congestive heart failure, 366–367, 371
screening for, 315

Sleepiness
definition of, 309
excessive daytime. *See also* Narcolepsy.
causes of, 310–311
diagnosis of, 309–312
differential diagnosis of, 337–338
drugs for, side effects of, 498–500
in kidney failure, 529
in obstructive sleep apnea, 337–338
measurement of, 309–310

Sleep-onset association disorder, 553

Sleep-onset REM period, in narcolepsy, 394–395, 404–405

Sleep-related abnormal swallowing syndrome, 425

Sleep-related painful erections, 425

Sleepwalking, 427–428
in pediatric patients, 553–554
overview of, 314–315

Small for gestational age, adult cardiovascular disease and, 973

Smoking
cessation of, for coronary artery disease prevention, 947–950, 964, 1073
obstructive sleep apnea in, 332

Smooth muscle relaxants, sleepiness due to, 504

Snapping hip syndromes, 196–197

Snoring
asthma exacerbation in, 520–521
description of, 329–330
in obstructive sleep apnea, 337, 569, 571

Social support, assessment of, 629

Sodium bicarbonate, as ergogenic aid, 287–288

Sodium intake, hypertension and, 726, 736, 1017–1018

Sodium oxybate (gamma hydroxybutyrate), for narcolepsy, 398, 400

Somnambulism. *See* Sleepwalking.

Somniloquy (sleep talking), 426, 554

Somnolence. *See* Sleepiness.

Sotalol, for atrial fibrillation, 1095, 1101

South Carolina Cardiovascular Disease Prevention (SCCDPP), 869, 872–873

Spear tackling, in football, cervical injury in, 233

Special Olympics, neck considerations in, 239–240

Speech therapy, progress in, review of, 634

Spilled teacup sign, in carpal instability, 61

Spinal stenosis
cervical injuries and, 235–236
lumbar injuries and, 215, 219–221

Spine
cervical, injuries of. *See* Cervical injuries.
pain related to. *See* Back pain.

Spiritual issues, in end-of-life care, 821–822

Spironolactone, for congestive heart failure, 1124

Splints
for bruxism, 423
for cubital tunnel syndrome, 10
for lateral epicondylitis, 5
for mallet finger, 97
for medial epicondylitis, 8
for plantar fasciitis, 156
for triceps tendon injury, 13

Spondylolisthesis, in athletes, 203, 208–210

Spondylolysis, in athletes, 204–208

Sports hernia, 178–179

Sports injuries
back pain in, **201–229**
biomechanics of, **245–251**
cervical pain in, **231–243**
elbow, **1–16**
ergogenic aids and, **277–292**
foot and ankle
examination of, **105–132**
types of, **133–161**
groin, **163–183, 185–200**
hand
anatomic considerations in, **71–89**
treatment of, **91–103**
osteochondral. *See* Osteochondral injuries.

wrist
anatomic considerations in, **17–33**
treatment of, **35–70**

Sprains
ankle. *See* Ankle injuries, sprains.
cervical, 237–238
finger, 86
first metatarsophalangeal joint, 156–157
Lisfranc joint, 126–127, 143–145
wrist, 58–62

Squeakers syndrome (intersection syndrome), 38–40

Squeeze test
in ankle sprains, 119
in Morton's neuroma, 128

Stable angina. *See* Angina, stable.

Standing, body alignment observation during, in foot and ankle injuries, 107–108

Stanford Five City Project, 867, 870–871

Stanford Three Community Study in California, 866–867, 870

Statins, for hyperlipidemia, 1045–1047

Stener lesion, 96–97

Steppage gait, in athletes, 109–110

Steroids, anabolic, as ergogenic aids, 280–282

Stimulants
as ergogenic aids, 279–280
for narcolepsy, 397–399
for sleep deprivation, 486–487

Stimulus control therapy, for insomnia, 381–382, 568–569

Stingers, in athletics, 234

Strains
adductor, 166–168
cervical, 237–238
high hamstring, 171
iliopsoas, 169–171
in back muscles, 213–214, 216
rectus femoris, 172
sartorius, 172

Strength testing, in hand injuries, 81–84

Strength training, for fall prevention, 692–694

Streptokinase, for coronary artery disease, 1071

Stress fractures
in spondylolysis, 204
of distal radius, 49
of femoral neck, 189–191
of foot, 146–148
of groin region, 189–191
of pubic ramus, 189–191
of sacrum, 220
of wrist, 36

Stress nuclear perfusion imaging, in coronary artery disease, 935–937

Stress testing, in coronary artery disease, 1060–1063, 1075
exercise electrocardiography in, 934–935, 940, 1060–1061
exercise nuclear perfusion imaging in, 936

Stress urinary incontinence, 708, 710

Stretching exercises, for fall prevention, 694

Stroke
central sleep apnea in, 367
hypertension causing, 1011
in atrial fibrillation, 1086–1087
obstructive sleep apnea and, 334–335
sleep disorders in, 540–541

Subluxation
of distal radioulnar joint, 44
of extensor carpi ulnaris, 41
of peroneal tendons, 154–155

Substance abuse, screening for, 633

Succinimides, sleepiness due to, 503

Sudden infant death syndrome, 552

Sudden unexplained nocturnal death syndrome, 428–429

Suicidal thoughts, in depression, 785–786

Sundowning, in Alzheimer's disease, 537

Supination, evaluation of, 111

Supination lift test, in triangular fibrocartilage complex injury, 44–45

Supplements, for sports performance. *See* Ergogenic aids.

Support groups, for dementia, 615

Supraventricular tachycardia, electrocardiography in, 905–910

Sural nerve, entrapment of, examination in, 124

Surgery
cardiac, atrial fibrillation after, 1096
for acetabular injury, 192–194

Surgery (*continued*)
for adductor strain, 168
for ankle sprain, 142–143
for back pain, 224–225
for biceps tendon injury, 12
for boxer's knuckle, 100
for carpal tunnel syndrome, 64–65
for cervical pain, 239
for cubital tunnel syndrome, 10–11
for DeQuervain's tenosynovitis, 38
for disc injury, 214, 218
for distal radius fracture, 48–49
for elbow rotatory instability, 11
for flexor carpi radialis tendonopathy, 42
for flexor carpi ulnaris tendonopathy, 42
for groin disruption, 180
for hamstring avulsion, 174
for hip avascular necrosis, 196
for iliopsoas disorders, 168
for jersey finger, 98–99
for Kienböck's disease, 56
for lateral femoral cutaneous nerve entrapment, 178
for Lisfranc's sprain, 145
for obstructive sleep apnea, 350–351
for obturator nerve entrapment, 175
for osteitis pubis, 188
for peroneal subluxation, 155
for posterior elbow impingement, 15
for rectus femoris strain, 173
for scaphoid fracture, 51–54
for skier's thumb, 96
for spondylolisthesis, 210
for sports hernia, 179
for talar osteochondral lesion, 153
for tarsal coalition, 152
for trapezium fracture, 58
for triangular fibrocartilage complex injury, 45
for triceps tendon injury, 13
for ulnar collateral ligament injury, 10
risks of, in obstructive sleep apnea, 336

Swallowing, sleep-related problems with, 425

Swan-neck deformity, of finger, 86

Swedish Trial in Old Patient with Hypertension, 739–740

Symphysis pubis
inflammation of, 186–188
osteomyelitis of, 188–189

Syndesmosis, ankle, sprains of, 142–143

Systemic lupus erythematosus, cardiovascular disease in, 971

Systolic Hypertension in Europe trial, 730–731, 738, 1013–1014

Systolic Hypertension in the Elderly Program (SHEP) trial, 730, 738, 742, 1013

T

T lymphocytes, sleep deprivation effects on, 480

T wave characteristics, in electrocardiography, 923

Tachyarrhythmias, in sick sinus syndrome, 905

Tachycardia
atrial, 907
atrioventricular nodal re-entry, 906
atrioventricular re-entry, 906–907
multifocal atrial, 906, 909–910
sinus, 906
supraventricular, 905–910
ventricular, 921–922

Tackling, spear, in football, cervical injury in, 233

Tailor's bunion, 128

Talar tilt test, in ankle sprain, 118–119, 136–137

Talocalcaneal coalition, 151–152

Talus, osteochondral injuries of, 121–122, 153, 260, 270, 272

Tangiers disease, 1042

Tarsal coalition, 151–152

Tarsal tunnel syndrome, 125
anterior, 126–127

Tau, in circadian clock, 449

Team approach
in end-of-life care, 814–815
in nursing facility care, 806–809

Technetium-99m, in perfusion imaging, in coronary artery disease, 935–936

Teeth, nocturnal grinding of (bruxism), 314, 421–424

Temazepam
for insomnia, 494, 496
potentially inappropriate prescription of, 765

Ten S's, for fall risk evaluation, 688–689

Tendonitis
Achilles tendon, 120–121

flexor hallucis longus, 125
iliopsoas, 169–171
peroneal, 124, 154

Tendonopathies, wrist, 37–42

Tenecteplase, for coronary artery disease, 1071

Tennis elbow, 4–5

Terminal illness, care during. *See* End-of-life care.

Terrors, sleep, 314–315, 426–427

Testosterone, as ergogenic aid, 280–282

Thallium-201, in perfusion imaging, in coronary artery disease, 935–936

Thenar eminence, palpation of, 77–78

Theophylline
for central sleep apnea, 370
for nocturnal asthma, 521

Thioperamide, for narcolepsy, 400

Thioridazine, potentially inappropriate prescription of, 768

Thompson test, in Achilles tendon injuries, 120–121

Thrombolytic agents, for coronary artery disease, 1067–1068, 1071

Throwing
cubital tunnel syndrome in, 10–11
posterior elbow impingement in, 13–15
ulnar collateral ligament injury in, 8–10

Thumb injuries
anatomic considerations in, 72–73
Bennett fracture, 102
De Quervain's tenosynovitis, 37–38, 95
history of, 73–74
mallet thumb, 97–98
physical examination in, 75–88
Finkelstein test in, 87
inspection in, 75, 77
neurovascular, 80–81
palpation in, 77–78
range of motion assessment in, 78–80
strength testing in, 82–83
skier's thumb, 87–88, 96–97
Stener lesion, 96–97

Thyroid, desiccated, potentially inappropriate prescription of, 768

Tibial nerve, injuries of, evaluation of, 115
anterior, 125
posterior, 123–125

Ticlopidine, potentially inappropriate prescription of, 767

Timed Get Up and Go Test, for fall risk evaluation, 690

Tinea pedis, 112, 114

Tinel's sign
in carpal tunnel syndrome, 64
in tarsal tunnel syndrome, 125
in wrist injuries, 28–29

Tioclopidine, for coronary artery disease, 1068

Tiotropium, for COPD, 524

Tiptoe test, in ankle injuries, 110

Tirofiban, for coronary artery disease, 1068

Toe(s), injuries of, 127–129, 156–157

Toenail disorders, in athletes, 114–115

Tongue, surgery on, for obstructive sleep apnea, 350

Tonsil hypertrophy, obstructive sleep apnea in, 557

"Too-many-toes" sign, in hindfoot injuries, 123

Torcetrapib, for hyperlipidemia, 1050

Torus fractures, of distal radius, 48

Transferritin, deficiency of, restless legs syndrome in, 439, 441

Transient urinary incontinence, 702–703

Transplantation
fetal tissue, for narcolepsy, 401
heart, hyperlipidemia in, 1052
kidney, hyperlipidemia in, 1052

Tranylcypromine, for insomnia, 499

Trapezium
anatomy of, 21
fracture of, 58

Trapezoid, anatomy of, 21

Trazodone
for depression, 787–788
for insomnia, 383
sleepiness due to, 498

Triamcinolone, for DeQuervain's tenosynovitis, 95

Triamterene, for hypertension, 738

Triangular fibrocartilage complex
anatomy of, 20–21
examination of, 27
injury of, 44–45

Triazolam
for insomnia, 384, 494, 496
potentially inappropriate prescription of, 765

Triceps tendon, at elbow, injury of, 13

Tricuspid valve, regurgitation of, 1110

Tricyclic antidepressants
for depression, 787–788
for insomnia, 498–499
for narcolepsy, 399–400
sleepiness due to, 503

Trifascicular block, electrocardiography in, 920

Trigger fingers and thumb, 78, 99

Trigger point injections, for back pain, 225

Triglycerides, excess of. *See* Hypertriglyceridemia.

Trimethobenzamide, potentially inappropriate prescription of, 764

Trimipramine, for insomnia, 499

Tripelennamine, potentially inappropriate prescription of, 766

Triquetrum
anatomy of, 20
fracture of, 58

Tryptophan, for insomnia, 498

Tumor necrosis factor-α, narcolepsy and, 397

Turf toe, 156–157

U

U wave characteristics, in electrocardiography, 923–924

Ulna, styloid process of, palpation of, 77

Ulnar artery
at elbow, anatomy of, 3
in hand, examination of, 81

Ulnar collateral ligament
at elbow, strain of, 8–10
injury of, skier's thumb in, 87–88, 96–97

Ulnar nerve
at elbow
anatomy of, 2
compression of, 10–11
at wrist
anatomy of, 24
compression of, 65
in hand, examination of, 80

Ulnar variance, in triangular fibrocartilage complex injury, 45

Ultradian rhythms, in sleep, 321

Ultrasonography
brachial artery, in endothelial function evaluation, 941–942
in adductor strains, 167

United Kingdom Prospective Diabetes studies, 1021

Unstable angina. *See* Angina, unstable.

Urge urinary incontinence, 708, 711

Urinary incontinence, **699–722**
age-related, 700–701
classification of, 708–709
diagnosis of, 708
epidemiology of, 699–700
evaluation of, 635, 701–703
functional, 713
history of, 703, 705–706
information resources for, 715
mixed, 708, 712
nocturnal, 700, 714
overflow, 712
patient education on, 709, 715
physical examination in, 706–708
referral in, 717
risk factors for, 703–704
stress, 708, 710
transient, 702–703
treatment of, 709–717
nonpharmacologic, 709–716
pharmacologic, 716–717
urge, 708, 711
voiding diary in, 705, 719

Urokinase, for coronary artery disease, 1071

Uvula, surgery on, for obstructive sleep apnea, 350

Uvulopalatopharyngoplasty, for obstructive sleep apnea, 350, 573

V

Vaccinations, for elderly persons, 606

Valerian, for insomnia, 498

Valproic acid, for depression, 787

Valsartan, for coronary artery disease, 1069

Valvular heart disease, **1109–1114**. *See also specific disorders.*

Vascular dementia, diagnostic criteria for, 838–839

Vascular depression, 782

Venlafaxine
for depression, 787–788
for insomnia, 499

Ventricle(s)
left, hypertrophy of
coronary artery disease risk in, 966
electrocardiography in, 915–916
right, hypertrophy of, 903, 916

Ventricular septal defect, murmurs in, 1110

Ventricular tachycardia,
electrocardiography in, 921–922

Verapamil
for atrial fibrillation, 1091–1092
for coronary artery disease, 1069

Vertebral fractures, in athletes, 204–208, 212–213, 215, 218–219

Vesper's curse, 440

Veterans Administration Cooperative Trial, for hypertension, 739

Vitamin B complex, for hyperhomocysteinemia, 968

Vitamin requirements, for elderly persons, 660, 665

Voiding diary, for urinary incontinence, 705, 719

Volar intercalated segment instability, 62

Volar plate injuries, of fingers, 86, 99

W

Wakefulness, physiology of, 535–536

Walkers, for fall prevention, 695–696

Walking, in sleep, 314–315, 427–428

Wandering atrial pacemaker (multifocal atrial tachycardia),
electrocardiography in, 909–910

Warfarin, for atrial fibrillation, 1099–1100

Wartenberg's syndrome, 65–66

Warts, plantar, 112

Watson's test, in wrist injuries, 29, 59

Weight loss
in end-of-life care, 819
in malnutrition, 664

Weight training, for fall prevention, 692–693

Wenckebach block, electrocardiography in, 912–914

Whiplash injury, in sports, 237

Windlass mechanism test, in plantar fasciitis, 125

Wolff-Parkinson-White syndrome
atrial fibrillation in, 1092
atrioventricular re-entry tachycardia in, 907
pre-excitation syndromes in, 911

Women's Health Initiative, hypertension studies of, 1020

World Health Organization, insomnia nosology of, 378–379

Wrist actigraphy
in delayed sleep phase circadian sleep disorders, 455
in sleep deprivation, 483–484

Wrist injuries, **17–33, 35–70**
anatomic considerations in
bony, 17–22
clinical, 17
soft tissue, 22–24
surface, 25–26
approach to, 36–37
carpal tunnel syndrome, 64–65
compressive neuropathies, 63–66
cyclist's palsy, 65
diagnosis of, 30–32
distal posterior interosseous nerve syndrome, 65
distal radioulnar joint, 42–44
epidemiology of, 35–36
fractures, 45–58
capitate, 58
Colles', 46, 48
epidemiology of, 36
hamate, 56–57
Kienböck's disease, 55–56
lunate, 55–56
pisiform, 57–58
radius, 45–49
scaphoid, 49–55
trapezium, 58
triquetral, 58
ganglia, 63
ligamentous, 58–62
overuse, epidemiology of, 36
physical examination in, 24–30, 93–95
tendonopathies, 37–42
DeQuervain's tenosynovitis, 37–38, 95
extensor carpi ulnaris, 40–41
flexor carpi radialis, 42
flexor carpi ulnaris, 42
intersection syndrome, 38–40

Wrist injuries (*continued*)
triangular fibrocartilage complex, 44–45
types of, 36
Wartenberg's syndrome, 65–66

X

Xanthoma, in familial hypercholesterolemia, 1004

Ximelagatran, for atrial fibrillation, 1100

Y

Zaleplon, for insomnia, 383–384, 494, 496

Zarit burden interview, for caregivers, 851–852

Zolpiclone, for insomnia, 383

Zolpidem, for insomnia, 383–384, 494–496